

Sources and Credits

The publisher gratefully acknowledges the generous authors and publishers who have granted permission to reprint or adapt their works for inclusion in this volume. For readers' ease in locating specific materials, the following sources and credits are listed by chapter, with text figures cited first, followed by boxed figures and then tables.

Part 1 opening photograph:

Galen Rowell/Mountain Light Photography Emeryville, CA.

Chapter 1

Figure 1.1B: Staehelin, L. A., Giddings, T. H. Jr., Kiss, J. Z., Sack, F. D. (1990) Macromolecular differentiation of Golgi stacks in root tips of *Arabidopsis* and *Nicotiana* seedlings as visualized in high-pressure frozen and freeze-substituted samples. *Protoplasma* 157: 75–91. Copyright © 1990 Springer-Verlag.

Figure 1.11: A. L. Samuels, University of British Columbia, Vancouver, Canada; previously unpublished.

Figure 1.12: Reproduced from Tilney, L., Cooke, T. J., Connelly, P. S., Tilney, M. S. (1991) The structure of plasmodesmata as revealed by plasmolysis, detergent extraction, and protease digestion. *J. Cell Biol.* 112: 739–748. By copyright permission of the Rockefeller University Press.

Figure 1.13A: Oparka, K., Prior, D.A.M., Crawford, J. W. (1994) Behaviour of plasma membrane, cortical ER and plasmodesmata during plasmolysis of onion epidermal cells. *Plant Cell Environ.* 17: 163–171. Used with permission of Blackwell Science Ltd.

Figure 1.16A,B: Knebel, W., Quader, H., Schnepf, E. (1990) Mobile and immobile endoplasmic reticulum in onion bulb epidermis cells: short-term and long-term observations with a confocal laser scanning microscope. *Eur. J. Cell Biol.* 52: 328–340. Used with permission from Urban & Fischer Verlag, Niederlassung Jena, Germany.

Figure 1.17A,B: Reproduced from Bonnett, H. T., Newcomb, E. H. (1965) Polyribosomes and cisternal accumulations in root cells of radish. *J. Cell Biol.* 27: 423–432. By copyright permission of the Rockefeller University Press.

Figure 1.18: G. W. Turner, Washington State University, Pullman; previously unpublished.

Figure 1.19: Lichtscheidl, I. K., Lancelle, S. A., Hepler, P. K. (1990) Actin–endoplasmic reticulum complexes in *Drosera*: their structural relationship with the plasmalemma, nucleus, and organelles in cells prepared by high pressure freezing. *Protoplasma* 155: 116–126. Copyright © 1990 Springer-Verlag.

Figure 1.20B: Fernandez, D., Staehelin, L. A. (1987) Does gibberellic acid induce the

transfer of lipase from protein bodies to lipid bodies in barley aleurone cells? *Plant Phys.* 85: 487–496. Copyright © 1988 Gustav Fischer Verlag.

Figure 1.21B: B. A. Larkins, University of Arizona, Tucson; previously unpublished.

Figure 1.22: Craig, S., Staehelin, L. A. (1988) High-pressure freezing of intact plant tissues: evaluation and characterization of novel features of the endoplasmic reticulum and associated membrane systems. *Eur. J. Cell Biol.* 46: 81–93. Used with permission from Urban & Fischer Verlag, Niederlassung Jena, Germany.

Figure 1.23B: Staehelin, L. A., Giddings, T. H. Jr., Kiss, J. Z., Sack, F. D. (1990) Macromolecular differentiation of Golgi stacks in root tips of *Arabidopsis* and *Nicotiana* seedlings as visualized in high-pressure frozen and freeze-substituted samples. *Protoplasma* 157: 75–91. Copyright © 1990 Springer-Verlag.

Figure 1.24: Craig, S., Staehelin, L. A. (1988) High-pressure freezing of intact plant tissues: evaluation and characterization of novel features of the endoplasmic reticulum and associated membrane systems. *Eur. J. Cell Biol.* 46: 81–93. Used with permission from Urban & Fischer Verlag, Niederlassung Jena, Germany.

Figure 1.25: A. Nebenfuhr, University of Colorado, Boulder; previously unpublished.

Figure 1.26: Craig, S., Staehelin, L. A. (1988) High-pressure freezing of intact plant tissues: evaluation and characterization of novel features of the endoplasmic reticulum and associated membrane systems. *Eur. J. Cell Biol.* 46: 81–93. Used with permission from Urban & Fischer Verlag, Niederlassung Jena, Germany.

Figure 1.27: Staehelin, L. A., Giddings, T. H. Jr., Kiss, J. Z., Sack, F. D. (1990) Macromolecular differentiation of Golgi stacks in root tips of *Arabidopsis* and *Nicotiana* seedlings as visualized in high-pressure frozen and freeze-substituted samples. *Protoplasma* 157: 75–91. Copyright © 1990 Springer-Verlag.

Figure 1.29A: Staehelin, L. A., Giddings, T. H. Jr., Kiss, J. Z., Sack, F. D. (1990) Macromolecular differentiation of Golgi stacks in root tips of *Arabidopsis* and *Nicotiana* seedlings as visualized in high-pressure frozen and freeze-substituted samples. *Protoplasma* 157: 75–91. Copyright © 1990 Springer-Verlag.

Figure 1.29B: Y. Mineyuki, Hiroshima University, Japan; previously unpublished.

Figure 1.30: Craig, S., Staehelin, L. A. (1988) High-pressure freezing of intact plant tissues: evaluation and characterization of novel features of the endoplasmic reticulum and associated membrane systems. *Eur. J. Cell Biol.* 46: 81–93. Used with permission from Urban & Fischer Verlag, Niederlassung Jena, Germany.

Figure 1.31A,B: Fowke, L. C., Tanchak, M. A., Galway, M. E. (1991) Ultrastructural cytology of the endocytotic pathway in plants. In *Endocytosis, Exocytosis and Vesicle Traffic in Plants*, C. R. Hawes, J.O.D. Coleman, and D. E. Evans, eds. Cambridge University Press, Cambridge, UK, pp. 15–40. Reprinted with the permission of Cambridge University Press.

Figure 1.32: L. A. Staehelin, University of Colorado, Boulder; previously unpublished.

Figure 1.33: L. A. Staehelin, University of Colorado, Boulder; previously unpublished.

Figure 1.34: P. C. Bethke, University of California, Berkeley; previously unpublished.

Figure 1.35: Micrograph: E. H. Newcomb, University of Wisconsin, Madison; previously unpublished.

Figure 1.36: L. A. Staehelin, University of Colorado, Boulder; previously unpublished.

Figure 1.37B: T. Murata, University of Tokyo, Japan; previously unpublished.

Figure 1.38: E. H. Newcomb, University of Wisconsin, Madison; previously unpublished.

Figure 1.39A,B: S. E. Frederick, Mount Holyoke College, South Hadley, MA; previously unpublished.

Figure 1.40A: Frederick, S. E., Newcomb, E. H. (1975) Plant microbodies. *Protoplasma* 84: 1–29. Copyright © 1975 Springer-Verlag.

Figure 1.41B: P. J. Gruber, Mt. Holyoke College, South Hadley, MA; previously unpublished.

Figure 1.43: R. H. Köhler, Cornell University, Ithaca, NY; previously unpublished.

Figure 1.45: E. H. Newcomb, University of Wisconsin, Madison; previously unpublished.

Figure 1.46: W. P. Wergin, Agricultural Research Service, Beltsville, MD; previously unpublished.

Figure 1.47: Source: Smith, J. D., Todd, P., Staehelin, L. A. (1997) Modulation of statolith mass and grouping in white clover *Trifolium repens* growth in 1-g, microgravity and on the Clinostat. *Plant J.* 12: 1361–1373. Used with permission from Blackwell Science Ltd.

Figure 1.48: G. W. Turner, Washington State University, Pullman; previously unpublished.

Figure 1.49A: S. E. Frederick, Mount Holyoke College, South Hadley, MA; previously unpublished.

Figure 1.50: L. A. Staehelin, University of Colorado, Boulder; previously unpublished.

Figure 1.51: W. P. Wergin, Agricultural Research Service, Beltsville, MD; previously unpublished.

Figure 1.52: W. P. Wergin, Agricultural Research Service, Beltsville, MD; previously unpublished.

Figure 1.53: W. P. Wergin, Agricultural Research Service, Beltsville, MD; previously unpublished.

Figure 1.54: L. A. Staehelin, University of Colorado, Boulder; previously unpublished.

Figure 1.56: L. A. Staehelin, University of Colorado, Boulder; previously unpublished.

Figure 1.57: P. J. Gruber, Mt. Holyoke College, South Hadley, MA; previously unpublished.

Figure 1.58: E. H. Newcomb, University of Wisconsin, Madison; previously unpublished.

Figure 1.61: M. E. Doohan, Texas Tech University, Lubbock; previously unpublished.

Figure 1.62: M. E. Doohan, Texas Tech University, Lubbock; previously unpublished.

Chapter 2

Figure 2.1: N. J. Stacey, John Innes Centre, Norwich, UK; previously unpublished.

Figure 2.2A–C: K. Findlay, John Innes Centre, Norwich, UK; previously unpublished.

Figure 2.2D: P. Linstead, John Innes Centre, Norwich, UK; previously unpublished.

Figure 2.3: M. Bush, Duke University, Durham, NC; previously unpublished.

Figure 2.4B: Ledbetter, M. C., Porter, K. R. (1970) *Introduction to the Fine Structure of Plant Cells*. Springer-Verlag, New York. Copyright © 1970 Springer-Verlag.

Figure 2.5A,B: Knox, J. P., Day, S., Roberts, K. (1989) A set of cell surface glycoproteins forms an early marker of cell position, but not cell type, in the root apical meristem of *Daucus carota* L. *Development* 106: 47–56. Used with permission from Company of Biologists Ltd.

Figure 2.5C: P. Linstead, John Innes Centre, Norwich, UK; previously unpublished.

Figure 2.6: R. L. Nicholson, Purdue University, West Lafayette, IN; previously unpublished.

Figure 2.7: R. Cotter, Plant Gene Expression Center, Albany, CA; previously unpublished.

Figure 2.11B: Reproduced from Mueller, S. C., Brown, R. M. Jr. (1980) Evidence for an intramembrane component associated with a cellulose microfibril-synthesizing complex in higher plants. *J. Cell Biol.* 84: 315–326. By copyright permission of the Rockefeller University Press.

Figure 2.11E: M. Jarvis, Glasgow University, Scotland; previously unpublished.

Figure 2.13: The relationship of the orders is adapted from Dahlgren, G. (1989) An updated angiosperm classification. *Bot. J. Linn. Soc.* 100: 197–204. Reprinted by permission of Academic Press. Copyright © 1989.

Figure 2.18B: Stafstrom, J. P., Staehelin, L. A. (1986) The role of carbohydrate in maintaining extensin in an extended conformation. *Plant Physiol.* 81: 242–246.

Figure 2.18C: Reprinted by permission from Condit, C. M., Meagher, R. B. (1986) A gene encoding a novel glycine-rich structural protein of petunia. *Nature* 323: 178–181. Copyright © 1986 Macmillan Magazines Ltd.

Figure 2.19A–D: Ye, Z.-H., Varner, J. E. (1991) Tissue-specific expression of cell wall proteins in developing soybean tissues. *Plant Cell* 3: 23–38.

Figure 2.20D: Gane, A. M., Clarke, A. E., Bacic, A. (1995) Localisation and expression of arabinogalactan-proteins in the ovaries of *Nicotiana glauca* Link and Otto. *Sex. Plant Reprod.* 8: 278–282. Copyright © 1995 Springer-Verlag.

Figure 2.22: Adapted from Iiyama, K., Lam, T.B.-T., Stone, B. A. (1994) Covalent cross-links in the cell wall. *Plant Physiol.* 104: 315–320.

Figure 2.23A: N. Carpita, Purdue University, West Lafayette, IN; previously unpublished.

Figure 2.26B: Gunning, B.E.S., Steer, M. W. (1996) *Plant Cell Biology: Structure and Function*. Jones and Bartlett, Sudbury, MA. Copyright © 1996 and used by permission from Gustav Fischer Verlag.

Figure 2.26C: Zhang, G. F., Staehelin, L. A. (1992) Functional compartmentation of the Golgi apparatus of plant cells: immunocytochemical analysis of high-pressure frozen and freeze-substituted sycamore maple suspension culture cells. *Plant Physiol.* 99: 1070–1083.

Figure 2.30: Adapted from Gibeaut, D. M., Carpita, N. C. (1993) Synthesis of (1→3), (1→4) β -D-glucan in the Golgi apparatus of maize coleoptiles. *Proc. Natl. Acad. Sci. USA* 90: 3850–3854. Copyright © 1993 National Academy of Sciences, U.S.A.

Figure 2.31A–C: Reproduced from Giddings, T. H. Jr., Brower, D. L., Staehelin, L. A. (1980) Visualization of particle complexes in the plasma membrane of *Micrasterias denticulata* associated with the formation of cellulose fibrils in primary and secondary cell walls. *J. Cell Biol.* 84: 327–339. By copyright permission of the Rockefeller University Press.

Figure 2.31D: Herth, W. (1983) Arrays of plasma-membrane “rosettes” involved in cellulose microfibril formation of *Spirogyra*. *Planta* 157: 347–356. Copyright © 1983 Springer-Verlag.

Figure 2.31E,F: Montezinos, D., Brown, M. Jr. (1976) Surface architecture of the plant cell: biogenesis of the cell wall, with special emphasis on the role of the plasma membrane in cellulose biosynthesis. *J. Supramol. Struct.* 5: 277–290. Reprinted by permission of Wiley-Liss, Inc., a subsidiary of John Wiley & Sons, Inc.

Figure 2.31G,H: Grimson, M. J., Haigler, C. H., Blanton, R. L. (1996) Cellulose microfibrils, cell motility, and plasma membrane protein organization change in parallel dur-

ing culmination in *Dictyostelium discoideum*. *J. Cell Sci.* 109: 3079–3087. Used with permission from Company of Biologists Ltd.

Figure 2.31I–K: Herth, W. (1985) Plasma-membrane rosettes involved in localized wall thickening during xylem vessel formation of *Lepidium sativum*. *Planta* 164: 12–21. Copyright © 1985 Springer-Verlag.

Figure 2.36: Adapted from Cosgrove, D. J. (1993) How do plant cell walls extend? *Plant Physiol.* 102: 1–6.

Figure 2.37: Adapted from McQueen-Mason, S., Durachko, D. M., Cosgrove, D. J. (1992) Two endogenous proteins that induce cell wall extension in plants. *Plant Cell* 4: 1425–1433.

Figure 2.39A,B: M. Bush, Duke University, Durham, NC; previously unpublished.

Figure 2.39C,D: Stacy, N. J., Roberts, K., Carpita, N. C., Wells, B., McCann, M. C. (1995) Dynamic changes in cell surface molecules are very early events in the differentiation of mesophyll cells from *Zinnia elegans* into tracheary elements. *Plant J.* 8: 891–906. Used with permission from Blackwell Science Ltd.

Figure 2.40A–C: Bowes, B. G. (1996) *A Colour Atlas of Plant Structure*. Manson Publishers, London. Copyright © 1996 Manson Publishers Ltd.

Figure 2.41A: Beasley, C. A. (1975) Developmental morphology of cotton flowers and seed as seen with the scanning electron microscope. *Am. J. Bot.* 62: 584–592. Used with permission from the Botanical Society of America, Inc. Copyright © 1975.

Figure 2.41B–D: Ledbetter, M. C., Porter, K. R. (1970) *Introduction to the Fine Structure of Plant Cells*. Springer-Verlag, New York. Copyright © 1970 Springer-Verlag.

Figure 2.42: Micrographs reprinted from Falconer, M. M., Seagull, R. W. (1985) Immunofluorescent and Calcofluor white staining of developing tracheary elements in *Zinnia elegans* suspension cultures. *Protoplasma* 125: 190–198. Copyright © 1985 Springer-Verlag.

Figure 2.43: Bowes, B. G. (1996) *A Colour Atlas of Plant Structure*. Manson Publishers, London. Copyright © 1996 Manson Publishers Ltd.

Figure 2.44A: Dharmawardhana, D P., Ellis, B. E., Carlson, J. E. (1992) Characterization of vascular lignification in *Arabidopsis thaliana*. *Can. J. Bot.* 70: 2238–2244. Used with permission from NRC Research Press. Copyright © 1992.

Figure 2.44B: Bowes, B. G. (1996) *A Colour Atlas of Plant Structure*. Manson Publishers, London. Copyright © 1996 Manson Publishers Ltd.

Figure 2.44C: Zhong, R., Taylor, J. J., Ye, Z.-H. (1997) Disruption of interfascicular fiber differentiation in an *Arabidopsis* mutant. *Plant Cell* 9: 2159–2170.

- Figure 2.44D,E: Chapple, C.C.S., Vogt, T., Ellis, B. E., Somerville, C. R. (1992) An *Arabidopsis* mutant defective in the general phenylpropanoid pathway. *Plant Cell* 4: 1413–1424.
- Figure 2.45A: D. DeMason, University of California, Riverside; previously unpublished.
- Figure 2.45B: M. Buckeridge, Instituto de Botanica, Secao de Fisiologia e Bioquimica Plantas, Sao Paulo, SP, Brasil; previously unpublished.
- Figure 2.46A,B: McCann, M. C., Stacey, N. J., Wilson, R., Roberts, K. (1993) Orientation of macromolecules in the walls of elongating carrot cells. *J. Cell Sci.* 106: 1347–1356. Used with permission from Company of Biologists Ltd.
- Figure 2.46C: J. Labavitch, University of California, Davis; previously unpublished.
- Figure 2.46D: D. Delmer, University of California, Davis; previously unpublished.
- Figure 2.46E: C. Haigler, Texas Tech University, Lubbock; previously unpublished.
- Box 2.1B: N. Carpita, Purdue University, West Lafayette, IN; previously unpublished.
- Box 2.1D: Carpita, N. C., Shea, E. M. (1989) Linkage structure of carbohydrates by gas chromatography–mass spectrometry (GC-MS) of partially methylated alditol acetates. In *Analysis of Carbohydrates by GLC and MS*, C. Biermann and G. D. McGinnis, eds. CRC Press, Boca Raton, FL, pp. 157–216. Copyright © 1989 CRC Press.
- Box 2.2D,E: W. York, University of Georgia, Athens; previously unpublished.
- Box 2.3: N. Carpita, Purdue University, West Lafayette, IN; previously unpublished.
- Box 2.4A: Hooke, R. (1664) *Micrographia*. The Council of the Royal Society of London for Improving Natural Knowledge, London. Reprinted by permission of the President and Council of the Royal Society.
- Box 2.4B: McCann, M. C., Wells, B., Roberts, K. (1990) Direct visualization of cross-links in the primary plant cell wall. *J. Cell Sci.* 96: 323–334. Used with permission from Company of Biologists Ltd.
- Box 2.4C: McCann, M. C., Roberts, K., Wilson, R. H., Gidley, M. J., Gibeaut, D. M., Kim, J.-B., Carpita, N. C. (1995) Old and new ways to probe plant cell-wall architecture. *Can. J. Bot.* 73 (Suppl. 1): S103–S113. Used with permission from NRC Research Press. Copyright © 1995.
- Box 2.4D: Ledbetter, M. C., Porter, K. R. (1970) *Introduction to the Fine Structure of Plant Cells*. Springer-Verlag, New York. Copyright © 1970 Springer-Verlag.
- Box 2.4E: B. Wells, John Innes Center, Norwich, UK; previously unpublished.
- Box 2.4F: V. Morris, Institute of Food Research, Norwich, UK; previously unpublished.
- Box 2.5: Table adapted from Knox, J. P. (1989) The use of antibodies to study the architecture and developmental regulation of plant cell walls. *Int. Rev. Cytol.* 171: 79–120. Reprinted by permission of Academic Press. Copyright © 1989.
- Box 2.5A: Steele, N. M., McCann, M. C., Roberts, K. (1997) Pectin modification in cell walls of ripening tomatoes occurs in distinct domains. *Plant Physiol.* 114: 373–381.
- Box 2.5B: McCann, M. C., Roberts, K., Wilson, R. H., Gidley, M. J., Gibeaut, D. M., Kim, J.-B., Carpita, N. C. (1995) Old and new ways to probe plant cell-wall architecture. *Can. J. Bot.* 73 (Suppl.): S103–S113. Used with permission from NRC Research Press. Copyright © 1995.
- Box 2.5C: Zhang, G. F., Staehelin, L. A. (1992) Functional compartmentation of the Golgi apparatus of plant cells: immunocytochemical analysis of high-pressure frozen and freeze-substituted sycamore maple suspension culture cells. *Plant Physiol.* 99: 1070–1083.
- Box 2.5D,E: M. C. McCann, John Innes Center, Norwich, UK; previously unpublished.
- ### Chapter 3
- Figure 3.2: Johansson, F., Sommarin, M., Larsson, C. (1994) Rapid purification of the plasma membrane H^+ -ATPase in its non-activated form using FPLC. *Physiol. Plant.* 92: 389–396. Copyright © 1994 Physiology International and published by Munksgaard International Publishers, Ltd., Copenhagen, Denmark.
- Figure 3.3: Reprinted from Junge, W., Lill, H., Engelbrecht, S. (1997) ATP synthase: an electrochemical transducer with rotatory mechanics. *Trends Biochem. Sci.* 22: 420–423. Copyright © 1997 with permission from Elsevier Science.
- Figure 3.4: Adapted from (1997) ATP generation by photosynthesis. In *Plant Biochemistry and Molecular Biology*, H.-W. Heldt. Oxford University Press, London. Used with permission from Oxford University Press.
- Figure 3.7A: DeWitt, N. D., Sussman, M. R. (1995) Immunocytological localization of an epitope-tagged plasma membrane proton pump (H^+ -ATPase) in phloem companion cells. *Plant Cell* 7: 2053–2067.
- Figure 3.7B,C: Harper, J. F., Manney, L., Sussman, M. R. (1994) The plasma membrane H^+ -ATPase gene family in *Arabidopsis*: genomic sequence of *AHA10* which is expressed primarily in developing seeds. *Mol. Gen. Genet.* 244: 572–587. Copyright © 1994 Springer-Verlag.
- Figure 3.11: Adapted with permission from Finbow, M. E., Harrison, M. A. (1997) The vacuolar H^+ -ATPase: a universal proton pump of eukaryotes. *Biochem. J.* 324: 697–712. Copyright © 1997 The Biochemical Society.
- Figure 3.14A,B: Data adapted from Lu, Y.-P., Li, Z.-S., Drozdowicz, Y. M., Hoertensteiner, S., Martinoia, E., Rea, P. A. (1998) AtMRP2, an *Arabidopsis* ATP binding cassette transporter able to transport glutathione *S*-conjugates and chlorophyll catabolites: functional comparisons with AtMRP1. *Plant Cell* 10: 267–282.
- Figure 3.17: Graphs: D. Sanders, University of York, UK; previously unpublished.
- Figure 3.19A,B: Stadler, R., Sauer, N. (1996) The *Arabidopsis thaliana* AtSUC2 gene is specifically expressed in companion cells. *Bot. Acta* 109: 299–306. Copyright © 1996 Georg Thieme Verlag, Stuttgart–New York.
- Figure 3.20A,B: Reprinted with permission from Kühn, C., Franceschi, V. R., Schulz, A., Lemoine, R., Frommer, W. B. (1997) Macromolecular trafficking indicated by localization and turnover of sucrose transporters in enucleate sieve elements. *Science* 275: 1298–1302. Copyright © 1997 American Association for the Advancement of Science.
- Figure 3.22: Kim, E. J., Kwak, J. M., Uozumi, N., Schroeder, J. I. (1998) AtKUP1: an *Arabidopsis* gene encoding high-affinity potassium transport activity. *Plant Cell* 10: 51–62.
- Figure 3.25A: Roblin, G. (1979) *Mimosa pudica*: a model for the study of the excitability in plants. *Biol. Rev.* 54: 135–153. Reprinted with the permission of Cambridge University Press.
- Figure 3.25B: Abe, T., Oda, K. (1976) Resting and action potentials of excitable cells in the main pulvinus of *Mimosa pudica*. *Plant Cell Physiol.* 17: 1343–1346. Reprinted with permission from the Japanese Society of Plant Physiologists.
- Figure 3.28: Adapted from Pantoja, O., Dainty, J., Blumwald, E. (1989) Ion channels in vacuoles from halophytes and glyco-phytes *FEBS Lett.* 255: 92–96. Copyright © 1989 with permission from Elsevier Science.
- Figure 3.29: Adapted from Pantoja, O., Dainty, J., Blumwald, E. (1989) Ion channels in vacuoles from halophytes and glyco-phytes *FEBS Lett.* 255: 92–96. Copyright © 1989 with permission from Elsevier Science.
- Figure 3.32: Roberts, S. K., Tester, M. (1997) A patch clamp study of Na^+ transport in maize roots. *J. Exp. Bot.* 48 (Special Issue): 431–440. Used with permission from Oxford University Press.
- Figure 3.33A: Nakamura, R. L., McKendree, W. L. Jr., Hirsch, R. E., Sedbrook, J. C., Gaber, R. F., Sussman, M. R. (1995) Expression of an *Arabidopsis* potassium channel gene in guard cells. *Plant Physiol.* 109: 371–374.
- Figure 3.33B: Lagarde, D., Basset, M., Lepetit, M., Conejero, G., Gaymard, R., Astruc, S., Grignon, C. (1996) Tissue-specific expression of *Arabidopsis* AKT1 gene is consistent with a role in K^+ nutrition. *Plant J.* 9: 195–203. Used with permission from Blackwell Science Ltd.

Figure 3.35: Adapted and used with permission from Hirsch, R. E., Lewis, B. D., Spalding, E. P., Sussman, M. R. (1998) A role for the AKT1 potassium channel in plant nutrition. *Science* 280: 918–921. Copyright © 1998 American Association for the Advancement of Science.

Figure 3.37: Reprinted by permission from Kreuzsch, A., Pfaffinger, P. J., Stevens, C. F., Choe, S. (1998) Crystal structure of the tetramerization domain of the *Shaker* potassium channel. *Nature* 392: 945–948. Copyright © 1998 Macmillan Magazines Ltd.

Figure 3.39: Czempinski, K., Zimmerman, S., Ehrhardt, T., Müller-Röber, B. (1997) New structure and function in plant K⁺ channels: KCO₁, an outward rectifier with a steep Ca²⁺ dependency. *EMBO J.* 16: 2565–2575. Used with permission from Oxford University Press.

Figure 3.40: Roberts, S. K., Tester, M. (1997) A patch clamp study of Na⁺ transport in maize roots. *J. Exp. Bot.* 48 (Special Issue): 431–440. Used with permission from Oxford University Press.

Figure 3.41A,B: Allen, G. J., Amtmann, A., Sanders, D. (1998) Calcium-dependent and calcium-independent K⁺ mobilization channels in *Vicia faba* guard cell vacuoles. *J. Exp. Bot.* 49 (Special Issue): 305–318. Used with permission from Oxford University Press.

Figure 3.42: Adapted from Bethke, P. G., Jones, G., Jones, R. L. (1995) Calcium and plant hormone action (Fig. 1). In *Plant Hormones—Physiology, Biochemistry, and Molecular Biology*, P. J. Davies, ed. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp. 298–317. Used with kind permission from Kluwer Academic Publishers.

Figure 3.43: Adapted from Piñeros, M., Tester, M. (1995) Characterization of a voltage-dependent Ca²⁺-selective channel from wheat roots. *Planta* 195: 478–488. Copyright © 1995 Springer-Verlag.

Figure 3.45: Bethke, P., Jones, R. L. (1994) Ca²⁺-calmodulin modulates ion channel activity in storage protein vacuoles of barley aleurone cells. *Plant Cell* 6: 277–285.

Figure 3.46: Schroeder, J. I., Keller, B. U. (1992) Two types of anion channel currents in guard cells with distinct voltage regulation. *Proc. Natl. Acad. Sci. USA* 89: 5052–5029. Used with permission.

Figure 3.47: Adapted from Cerana, R., Giromini, L., Colombo, R. (1995) Malate-regulated channels permeable to anions in vacuoles of *Arabidopsis thaliana*. *Aust. J. Plant Physiol.* 22: 115–121. Used with permission from Commonwealth Scientific and Industrial Research Organization, Melbourne.

Figure 3.53: Reprinted by permission from Walz, T., Hirai, T., Murata, K., Heymann, J. B., Mitsuoka, K., Fujiyoshi, Y., Smith, B. L., Agre, P., Engel, A. (1997) The three-dimensional structure of aquaporin-1. *Nature* 387:

624–627. Copyright © 1997 Macmillan Magazines Ltd.

Box 3.1: D. Sanders, University of York, UK; previously unpublished.

Box 3.2A–C: Adapted from Uozumi, N., Tatsunosuke, N., Schroeder, J. I., Muto, S. (1998) Determination of transmembrane topology of an inward-rectifying potassium channel from *Arabidopsis thaliana* based on functional expression in *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 95: 9773–9778. Copyright © 1998 National Academy of Sciences, U.S.A.

Box 3.6: From product literature for the GeneClamp 500. Axon Instruments Inc., Foster City, CA.

Box 3.7B,C: Patch clamp tower: Used with permission from Adams & List Associates Ltd., Great Neck, NY.

Box 3.8A,B: Roberts, S. K., Tester, M. (1997) Permeation of Ca²⁺ and monovalent cations through an outwardly rectifying channel in maize root stelar cells. *J. Exp. Bot.* 48: 839–846. Used with permission from Oxford University Press.

Chapter 4

Figure 4.5B: K. Keegstra, Michigan State University, East Lansing; previously unpublished.

Figure 4.8A–C: M. Lee, R. Trelease, Arizona State University, Tempe; previously unpublished.

Figure 4.10A: A. Heese, N. Raikhel, Michigan State University, East Lansing; previously unpublished.

Figure 4.11A,B: Varagona, M. J., Schmidt, R. J., Raikhel, N. V. (1992) Nuclear localization signal(s) required for nuclear targeting of the maize regulatory protein Opaque-2. *Plant Cell* 4: 1213–1227.

Figure 4.11C,D: Hicks, G. R., Smith, H. M., Lobreaux, S., Raikhel, N. V. (1996) Nuclear import in permeabilized protoplasts from higher plants has unique features. *Plant Cell* 8:1337–1352.

Figure 4.23A,B: Reproduced from Lev-anony, J., Rubin, R., Altschuler, Y., Galili, G. (1992) Evidence for a novel route of wheat storage proteins to vacuoles. *J. Cell Biol.* 119: 1117–1128. By copyright permission of the Rockefeller University Press.

Figure 4.25: J. Dombrowski, N. Raikhel, Michigan State University, East Lansing; previously unpublished.

Figure 4.26: Bassham, D. C., Gal, S., Concaícao, A. D., Raikhel, N. V. (1995) An *Arabidopsis* syntaxin homologue isolated by functional complementation of a yeast pep12 mutant. *Proc. Natl. Acad. Sci. USA* 92: 7262–7266. Copyright © 1995 National Academy of Sciences, U.S.A.

Figure 4.31: D. Robinson, University of Goettingen, Germany; previously unpublished.

Box 4.1: Photomicrograph from Herman, E. M., Tague, B. W., Hoffman, L. M., Kjembrup, S. E., Chrispeels, M. J. (1990) Retention of phytohemagglutinin with carboxy-terminal tetrapeptide KDEL in the nuclear envelope and the endoplasmic reticulum. *Planta* 182: 305–312. Copyright © 1990 Springer-Verlag.

Box 4.4A–D: D. Robinson, University of Goettingen, Germany; previously unpublished.

Chapter 5

Figure 5.1: T. Baskin, University of Missouri, Columbia; previously unpublished.

Figure 5.2: Xu, P., Lloyd, C. W., Staiger, C. J., Drøbak, B. K. (1992) Association of phosphatidylinositol 4-kinase with the plant cytoskeleton. *Plant Cell* 4: 941–951.

Figure 5.6A: Reprinted by permission from Aebi, U., Cohn, J., Buhle, L., Gerace, L. (1986) The nuclear lamina is a meshwork of intermediate-type filaments. *Nature* 323: 560–564. Copyright © 1986 Macmillan Magazines Ltd.

Figure 5.7: Mizuno, K. (1995) A cytoskeletal 50kDa protein in higher plants that forms intermediate-sized filaments and stabilizes microtubules. *Protoplasma* 186: 99–112. Copyright © 1995 Springer-Verlag.

Figure 5.9A–F: Ross, J.H.E., Hutchings, A., Butcher, G. W., Lane, E. B., Lloyd, C. W. (1991) The intermediate filament-related system of higher plant cells shares an epitope with cytokeratin 8. *J. Cell Sci.* 99: 91–98. Used with permission from Company of Biologists Ltd.

Figure 5.10: Adapted from Linder, S., Schliwa, M., Kube-Granderath, E. (1997) Sequence analysis and immunofluorescence study of alpha- and beta-tubulins in *Reticulomyxa filosa*: implications of the high degree of beta-2-tubulin divergence. *Cell Motil. Cytoskeleton* 36: 164–178. Reprinted by permission of Wiley-Liss, Inc., a subsidiary of John Wiley & Sons, Inc.

Figure 5.11A: Adapted from Villemur, R., Joyce, C. M., Haas, N. A., Goddard, R. H., Kopczak, S. D., Hussey, P. J., Snustad, D. P., Silflow, C. D. (1992) α -Tubulin gene family of maize (*Zea mays* L.): evidence for two ancient α -tubulin genes in plants. *J. Mol. Biol.* 227: 81–96. Reprinted by permission of Academic Press. Copyright © 1992.

Figure 5.11B: Adapted from Villemur, R., Haas, N. A., Joyce, C. M., Snustad, D. P., Silflow, C. D. (1994) Characterization of four new beta-tubulin genes and their expression during male flower development in maize (*Zea mays* L.). *Plant Mol. Biol.* 24: 295–315. Used with kind permission from Kluwer Academic Publishers.

Figure 5.12A,B: Reproduced from Hoyle, H. D., Raff, E. C. (1990) Two *Drosophila* beta tubulin isoforms are not functionally equivalent. *J. Cell Biol.* 111: 1009–1026. By

copyright permission of the Rockefeller University Press.

Figure 5.13A,B: Carpenter, J. L., Ploense, S. E., Snustad, D. P., Silflow, C. D. (1992) Preferential expression of an α tubulin gene of *Arabidopsis* in pollen. *Plant Cell* 4: 557–571.

Figure 5.16A: Reprinted by permission from Kabsch, W., Mannherz, H. G., Suck, D., Pai, E. F., Holmes, K. C. (1990) Atomic structure of the actin DNase I complex. *Nature* 347: 37–44. Copyright © 1990 Macmillan Magazines Ltd.

Figure 5.16C: Reprinted by permission from Holmes, K. C., Popp, D., Gebhard, W., Kabsch, W. (1990) Atomic model of the actin filament. *Nature* 347: 44–49. Copyright © 1990 Macmillan Magazines Ltd.

Figure 5.16D: Reprinted by permission from Milligan, R. A., Whittaker, M., Safer, D. (1990) Molecular structure of F-actin and location of surface binding sites. *Nature* 348: 217–221. Copyright © 1990 Macmillan Magazines Ltd.

Figure 5.17A: Reprinted by permission from Nogales, E., Wolf, S. G., Downing, K. H. (1998) Structure of the alpha-beta tubulin dimer by electron crystallography. *Nature* 391: 199–203. Copyright © 1998 Macmillan Magazines Ltd.

Figure 5.17C: Juniper, B. E., Lawton, J. R. (1979) The effect of caffeine, different fixation regimes and low temperature on microtubules in the cells of higher plants. *Planta* 145: 411–416. Copyright © 1979 Springer-Verlag.

Figure 5.17D: D. Chrétien, EMBO Laboratory, Heidelberg, Germany; previously unpublished.

Figure 5.19B,D: Reproduced from Mandelkow, E.-M., Mandelkow, E., Milligan, R. A. (1991) Microtubule dynamics and microtubule caps: a time-resolved cryo-electron microscopy study. *J. Cell Biol.* 114: 977–992. By copyright permission of the Rockefeller University Press.

Figure 5.20A–D: Reproduced by permission from Hollenbeck, P. J. (1989) Dissecting a molecular motor. *Nature* 338: 294–295. Copyright © 1989 Macmillan Magazines Ltd.

Figure 5.24B,C: Williamson, R. E. (1975) Cytoplasmic streaming in *Chara*: a cell model activated by ATP and inhibited by cytochalasin B. *J. Cell Sci.* 17: 655–668. Used with permission from Company of Biologists Ltd.

Figure 5.24D: G. Wasteneys, Australian National University, Canberra; previously unpublished.

Figure 5.25A,B: Trojan, A., Gabryś, H. (1996) Chloroplast distribution in *Arabidopsis thaliana* (L.) depends on light conditions during growth. *Plant Physiol.* 111: 419–425.

Figure 5.26B: Mineyuki, Y., Kataoka, H., Masuda, Y., Nagai, R. (1995) Dynamic changes in the actin cytoskeleton during

the high-fluence rate response of the *Mougeotia* chloroplast. *Protoplasma* 185: 222–229. Copyright © 1995 Springer-Verlag.

Figure 5.27A,B: Lancelle, S. A., Cresti, M., Hepler, P. K. (1996) Growth inhibition and recovery in freeze-substituted *Lilium longiflorum* pollen tubes: structural effects of caffeine. *Protoplasma* 196: 21–33. Copyright © 1996 Springer-Verlag.

Figure 5.28A: Baskin, T. I., Miller, D. D., Vos, J. W., Wilson, J. E., Hepler, P. K. (1996) Cryofixing single cells and multicellular specimens enhances structure and immunocytochemistry for light microscopy. *J. Microsc.* 182: 149–161. Used with permission from Blackwell Science Ltd.

Figure 5.28B: Reprinted from Gunning, B.E.S., Hardham, A. R. (1979) Microtubules and morphogenesis in plants. *Endeavor* 3:112–117. Copyright © 1979 with permission from Elsevier Science.

Figure 5.29: Reproduced from Euteneuer, U., Jackson, W. T., McIntosh, J. R. (1982) Polarity of spindle microtubules in *Haemaphysalis* endosperm. *J. Cell Biol.* 94: 644–653. By copyright permission of the Rockefeller University Press.

Figure 5.30A: McDonald, A. R., Liu, B., Joshi, H. C., Palevitz, B. A. (1993) Gamma tubulin is associated with a cortical microtubule-organizing zone in the developing guard cells of *Allium cepa* L. *Planta* 191: 357–361. Copyright © 1993 Springer-Verlag.

Figure 5.30B,C: Liu, B., Joshi, H. C., Palevitz, B. A. (1995) Experimental manipulation of gamma-tubulin distribution in *Arabidopsis* using anti-microtubule drugs. *Cell Motil. Cytoskeleton* 31: 113–129. Reprinted by permission of Wiley-Liss, Inc., a subsidiary of John Wiley & Sons, Inc.

Figure 5.31A,B: Baskin, T. I., Wilson, J. E. (1997) Inhibitors of protein kinases and phosphatases alter root morphology and disorganize cortical microtubules. *Plant Physiol.* 113: 493–502.

Figure 5.32: Vesik, P. A., Vesik, M., Gunning, B.E.S. (1996) Field emission scanning electron microscopy of microtubule arrays in higher plant cells. *Protoplasma* 195: 168–182. Copyright © 1996 Springer-Verlag.

Figure 5.34: Zhang, D., Wadsworth, P., Hepler, P. K. (1990) Microtubule dynamics in living dividing plant cells: confocal imaging of microinjected fluorescent brain tubulin. *Proc. Natl. Acad. Sci. USA* 87: 8820–8824. Used with permission.

Figure 5.35A: Lloyd, C. W., Shaw, P. J., Warn, R. M., Yuan, M. (1996) Gibberellic acid induced reorientation of cortical microtubules in living plant cells. *J. Microsc.* 181: 140–144. Used with permission from Blackwell Science Ltd.

Figure 5.35B: Cleary, A.L. (1995) F-actin redistributions at the division site in living *Tradescantia* stomatal complexes as revealed by microinjection of rhodamine-phalloidin.

Protoplasma 185: 152–165. Copyright © 1995 Springer-Verlag.

Figure 5.36: R. Cyr, Pennsylvania State University, Abington; previously unpublished.

Figure 5.37: Adapted from Williamson, R. E. (1990) Alignment of cortical microtubules by anisotropic wall stresses. *Aust. J. Plant Physiol.* 17: 601–614 Used with permission from Commonwealth Scientific and Industrial Research Organization, Melbourne.

Figure 5.38: Vesik, P. A., Vesik, M., Gunning, B.E.S. (1996) Field emission scanning electron microscopy of microtubule arrays in higher plant cells. *Protoplasma* 195: 168–182. Copyright © 1996 Springer-Verlag.

Figure 5.39C: Reproduced from Waters, J. C., Chen, R.-H., Murray, A. W., Salmon, E. D. (1998) Localization of Mad2 to kinetochores depends on microtubule attachment, not tension. *J. Cell Biol.* 141: 1181–1191. By copyright permission of the Rockefeller University Press.

Figure 5.39D: Reproduced from Khodjakov, A., Cole, R. W., Bajer, A. S., Rieder, C. L. (1996) The force for poleward chromosome motion in *Haemaphysalis* cells acts along the length of the chromosome during metaphase but only at the kinetochore during anaphase. *J. Cell Biol.* 132: 1093–1104. By copyright permission of the Rockefeller University Press.

Figure 5.40C: Reproduced from Moritz, M., Braunfeld, M. B., Fung, J. C., Sedat, J. W., Alberts, B. M., Agard, D. A. (1995) Three-dimensional structural characterization of centrosomes from early *Drosophila* embryos. *J. Cell Biol.* 130: 1149–1159. By copyright permission of the Rockefeller University Press.

Figure 5.40D: H. Schatten, University of Missouri, Columbia, previously unpublished.

Figure 5.43A–K: Gunning, B.E.S., Steer, M. W. (1996) *Plant Cell Biology: Structure and Function*. Jones and Bartlett, Sudbury, MA. Copyright © 1996 and used by permission from Gustav Fischer Verlag.

Figure 5.44A: R. Martin, Gerhard Wanner, University of Munich, Germany; previously unpublished.

Figure 5.44C: Van Hooser, A., Brinkley, B. R. (1999) Methods for in situ localization of proteins and DNA in the centromere-kinetochore complex. *Methods Cell Biol.* 61: 57–80. Reprinted by permission of Academic Press. Copyright © 1999.

Figure 5.44D: K. McDonald, University of California, Berkeley; previously unpublished.

Figure 5.52A: Galatis, B., Apostolakis, P., Katsaros, C. (1984) Experimental studies on the function of the cortical cytoplasmic zone of the preprophase microtubule band. *Protoplasma* 122: 11–26. Copyright © 1984 Springer-Verlag.

Figure 5.52B,C: Kakimoto, T., Shibaoka, H. (1987) Actin filaments and microtubules in the preprophase band and phragmoplast of tobacco cells. *Protoplasma* 140: 151–156. Copyright © 1987 Springer-Verlag.

Figure 5.53A–F: Cleary, A. L., Gunning, B.E.S., Wasteneys, G. O., Hepler, P. K. (1992) Microtubule and F actin dynamics at the division site in living *Tradescantia* stamen hair cells. *J. Cell Sci.* 103: 977–988. Used with permission from Company of Biologists Ltd.

Figure 5.54: Mineyuki, Y., Gunning, B.E.S. (1990) A role for preprophase bands of microtubules in maturation of new cell walls, and a general proposal on the function of preprophase band sites in cell division in higher plants. *J. Cell Sci.* 97: 527–538. Used with permission from Company of Biologists Ltd.

Figure 5.56A–D: Gunning, B.E.S., Steer, M. W. (1996) *Plant Cell Biology: Structure and Function* Jones and Bartlett, Sudbury, MA. Copyright © 1996 and used by permission from Gustav Fischer Verlag.

Figure 5.56E,F: Smirnova, E. A., Cox, D. L., Bajer, A. S. (1995) Antibody against phosphorylated proteins (MPM-2) recognizes mitotic microtubules in endosperm cells of higher plant *Haemanthus*. *Cell Motil. Cytoskeleton* 31: 34–44. Reprinted with permission from Wiley-Liss, Inc., a subsidiary of John Wiley & Sons, Inc.

Figure 5.57A–V: Zhang, D., Wadsworth, P., Hepler, P. K. (1993) Dynamics of microfilaments are similar but distinct from microtubules during cytokinesis in living dividing plant cells. *Cell Motil. Cytoskeleton* 24: 151–155. Reprinted with permission from Wiley-Liss, Inc., a subsidiary of John Wiley & Sons, Inc.

Figure 5.58: Kakimoto, T., Shibaoka, H. (1988) Cytoskeletal ultrastructure of phragmoplast-nuclei complexes isolated from cultured tobacco cells. *Protoplasma* (Suppl. 2): 95–103. Copyright © 1988 Springer-Verlag.

Figure 5.59A–C: Venverloo, C. J., Libbenga, K. R. (1981) Cell division in *Nautilocalyx* explants. II Duration of cytokinesis and velocity of cell-plate growth in large highly vacuolate cells. *Z. Pflanzenphysiol.* 102: 389–395. Used with permission from Urban & Fischer Verlag, Niederlassung Jena, Germany.

Figure 5.59D: Asada, T., Shibaoka, H. (1994) Isolation of polypeptides with microtubule-translocating activity from phragmoplasts of tobacco BY-2 cells. *J. Cell Sci.* 107: 2249–2257. Used with permission from Company of Biologists Ltd.

Figure 5.60A: Asada, T., Shibaoka, H. (1994) Isolation of polypeptides with microtubule-translocating activity from phragmoplasts of tobacco BY-2 cells. *J. Cell Sci.* 107: 2249–2257. Used with permission from Company of Biologists Ltd.

Figure 5.60B. Asada, Osaka University, Japan, previously unpublished.

Figure 5.61A: Gunning, B.E.S., Steer, M. W. (1996) *Plant Cell Biology: Structure and Function*. Jones and Bartlett, Sudbury, MA. Copyright © 1996 Gustav Fischer Verlag.

Figure 5.61B: Kakimoto, T., Shibaoka, H. (1992) Synthesis of polysaccharides in phragmoplasts isolated from tobacco BY-2 cells. *Plant Cell Physiol.* 33: 353–361. Used with permission from the Japanese Society of Plant Physiology.

Box 5.1A: Portrait used by permission of the Linnean Society of London.

Box 5.1B,C: B. J. Ford, Rothay House, Cambridgeshire, UK; previously unpublished.

Box 5.2: Ribbon diagrams adapted from Federov, A. A., Ball, T., Mahoney, N. M., Valenta, R., Almo, S. C. (1997) Molecular basis for allergen cross-reactivity: crystal structure and IgE-epitope mapping of birch pollen profilin. *Structure* 5: 33–45. Used with permission from Current Biology Ltd.

Box 5.2A,B: Valster, A. H., Pierson, E. S., Valenta, R., Hepler, P. K., Emons, A.M.C. (1997) Probing the plant actin cytoskeleton during cytokinesis and interphase by profilin microinjection. *Plant Cell* 9: 815–824.

Box 5.2C: Upper panel: Miller, D. D., Lancelle, S. A., Hepler, P. K. (1996) Actin microfilaments do not form a dense meshwork in *Lilium longiflorum* pollen tubes. *Protoplasma* 195:123–132. Copyright © 1996 Springer-Verlag. Middle and lower panels: Vidali, L., Hepler, P. K. (1997) Characterization and localization of profilin pollen grains and tubes of *Lilium longiflorum*. *Cell Motil. & Cytoskeleton* 36:323–338. Reprinted with permission from Wiley-Liss, Inc., a subsidiary of John Wiley & Sons, Inc.

Box 5.3A: Inoué, S. (1953) Polarization optical studies of the mitotic spindle. I The demonstration of spindle fibers in living cells. *Chromosoma* 5: 487–506. Copyright © 1953 Springer-Verlag.

Box 5.3B: Inoué, S., Bajer, A. (1961) Birefringence in endosperm mitosis. *Chromosoma* 12: 48–63. Copyright © 1961 Springer-Verlag.

Box 5.3C: Oldenbourg, R., Mei, G. (1995) New polarized light microscope. *J. Microsc.* 180: 140–147. Used by permission from Blackwell Science Ltd.

Box 5.4A–D: Kiss, J. Z., Giddings, T. H. Jr., Staehelin, L. A., Sack, F. D. (1990) Comparison of the ultrastructure of conventionally fixed and high-pressure frozen/freezesubstituted root tips of *Nicotiana* and *Arabidopsis*. *Protoplasma* 157: 64–74. Copyright © 1990 Springer-Verlag.

Box 5.4E,F: Reproduced (E) and adapted (F) from Samuels, A. L., Giddings, T. H., Staehelin, L. A. (1995) Cytokinesis in tobacco BY-2 and root tip cells: a new model of cell plate formation in higher plants. *J. Cell Biol.* 130: 1345–1357. By copyright permission of the Rockefeller University Press.

Part 2 opening photograph:

Galen Rowell/Mountain Light Photography, Emeryville, CA.

Chapter 6

Chapter 6 opening photograph: D. Richardson, Northampton, MA; published in Purves, W., Orians, G., Heller, H. C., Sadava, D. (1998) *Life: The Science of Biology*, 5th edition. Sinauer/Freeman, Sunderland, MA, p. 65. Used with permission from the photographer.

Figure 6.10B: Reprinted from Shippen, D. I., McKnight, T. (1998) Telomeres, telomerase, and plant development. *Trends Plant Sci.* 3: 126–129. Copyright © 1998 with permission from Elsevier Science.

Figures 6.12–6.16: Adapted from Cooper, G. M. (1997) *The Cell: A Molecular Approach*. Copublished by ASM Press, Washington, DC, and Sinauer Associates, Sunderland, MA.

Figure 6.21: Adapted from Cooper, G. M. (1997) *The Cell: A Molecular Approach*. Copublished by ASM Press, Washington, DC, and Sinauer Associates, Sunderland, MA.

Figure 6.22: Photomicrographs: Madigan, M. T., Martinko, J. M., Parker, J. (1997) *Brock Biology of Microorganisms*, 8th ed. Prentice Hall, Upper Saddle River, NJ.

Figure 6.23 top: Hiratsuka, J., Shimada, H., Whittier, R., Ishibashi, T., Sakamoto, M., Mori, M., Kondo, C., Honji, Y., Sun, C.-R., Meng, B.-Y., Li, Y.-Q., Kanno, A., Nishizawa, Y., Hirai, A., Shinozaki, K., Sugiura, M. (1989) The complete sequence of the rice *Oryza sativa* chloroplast genome: intermolecular recombination between distinct transfer RNA genes accounts for a major plastid DNA inversion during the evolution of the cereals. *Mol. Genet.* 217: 185–194. Copyright © 1989 Springer-Verlag.

Figure 6.23 bottom: Adapted from Maier, L.R.M., Neckermann, K., Igloi, G. L., Kössel, H. (1995) Complete sequence of the maize chloroplast genome: gene content, hotspots of divergence and fine tuning of genetic information by transcription editing. *J. Mol. Biol.* 251: 614–628. Adapted by permission of Academic Press. Copyright © 1995.

Figure 6.25: Adapted from Kornberg, A., Baker, T. A. (1992) *DNA Replication*, 2nd ed. W. H. Freeman, New York. Used with permission.

Figure 6.26: Fauron, C., Casper, M., Gao, Y., Moore, B. (1995) The maize mitochondrial genome: dynamic, yet functional. *Trends Genet.* 11: 228–235. Copyright © 1995 with permission from Elsevier Science.

Figure 6.27A–C: Fauron, C., Casper, M., Gao, Y., Moore, B. (1995) The maize mitochondrial genome: dynamic, yet functional. *Trends Genet.* 11: 228–235. Copyright © 1995 with permission from Elsevier Science.

Figure 6.28: DNA diagram adapted from Boer, P. H., Gray, M. W. (1991) Short

dispersed repeats localized in spacer regions of *Chlamydomonas reinhardtii* mitochondrial DNA. *Curr. Genet.* 19: 309–312.

Figure 6.29: Cover of *Plant Mol. Biol. Rep.* 11 (June), accompanying article by Hirai, A., Nakasono, M. (1993). Used with permission from the International Society for Plant Molecular Biology.

Figure 6.30: Bacterial structure adapted from Cooper, G. M. (1997) *The Cell: A Molecular Approach*. Copublished by ASM Press, Washington, DC, and Sinauer Associates, Sunderland, MA.

Figure 6.33: Graph: M. Sugiura, Center for Gene Research, Nagoya University, Japan; previously unpublished.

Figure 6.35: Adapted from Harris, E. H., Boynton, J. E., Gillham, N. W. (1994) Chloroplast ribosomes and protein synthesis. *Microbiol. Rev.* 58: 700–754. Used with permission from the American Society for Microbiology.

Figure 6.36: Adapted from Dudock, B. S., Katz, G., Taylor, E. K., Holley, R. (1969) Primary structure of wheat germ phenylalanine transfer RNA. *Proc. Natl. Acad. Sci. USA* 62: 941–945. Used with permission.

Figure 6.37: Adapted from Canaday, J., Guillemaut, P., Weil, J. H. (1980) The nucleotide sequences of the initiator transfer RNAs from bean cytoplasm and chloroplasts. *Nucleic Acids Res.* 8: 999–1008. Copyright © 1980 Springer-Verlag.

Figure 6.42: Adapted from Westaway, S. K. (1995) Splicing of tRNA precursors. In *tRNA: Structure, Biosynthesis, and Function*, D. Soell and U. L. Rajbhandary, eds. ASM Press, Washington, DC, pp. 79–92. Used with permission from the American Society for Microbiology.

Figure 6.43: Adapted from Cooper, G. M. (1997) *The Cell: A Molecular Approach*. Copublished by ASM Press, Washington, DC, and Sinauer Associates, Sunderland, MA.

Figure 6.46: Adapted from Westhoff, P., Herrmann, R. G. (1988) Complex RNA maturation in chloroplasts: the Psb-B operon from spinach. *Eur. J. Biochem.* 171: 551–564. Used with permission from Blackwell Science Ltd.

Figure 6.47: Adapted from Russell, P. (1998) *Genetics*, 5th ed. Addison Wesley Longman Publishers, Menlo Park, CA. Reprinted by permission of Addison Wesley Educational Publishers, Inc.

Box 6.2: Photograph: M. Sugiura, Center for Gene Research, Nagoya University, Japan; previously unpublished.

Box 6.4: W. Gruissem, University of California, Berkeley; previously unpublished.

Chapter 7

Figure 7.1: Rattner, J. B. 1991. The structure of the mammalian centromere. *BioEssays* 13: 51–56. Used with permission from Wiley-

Liss, Inc., a subsidiary of John Wiley & Sons, Inc.

Figure 7.9: L. Jesaitis, University of California, Berkeley; previously unpublished.

Figure 7.10A,B: Rhoades, M. M. (1950) Meiosis in maize. *J. Hered.* 41: 58–67. Used with permission from Oxford University Press.

Figure 7.11: Photographs from F. W. Goro, published in Federov, N. V. (1984) Transposable genetic elements in maize. *Sci. Am.* 250(6): 84–98. Used with permission of the Estate of F. W. Goro.

Figure 7.12: A. Brousseau, St. Mary's College, Marga, CA. Copyright © 1995 Brother Eric Vogel, St. Mary's College.

Figure 7.20: M. P. Tolson, University of California, Davis; previously unpublished.

Figure 7.22A: Meyne, J. (1993) In *Chromosomes: A Synthesis*, R. P. Wagner, M. P. Maguire, R. L. Stallings, eds. Wiley-Liss, New York. Used with permission from Wiley-Liss, Inc., a subsidiary of John Wiley & Sons, Inc.

Figure 7.22B: Reprinted from Willard, H. F. (1990) Centromeres of mammalian chromosomes. Reprinted from *Trends Genet.* 6: 410–416. Copyright © 1990 with permission from Elsevier Science.

Figure 7.22C: S. Ruzin, University of California, Berkeley; previously unpublished.

Figure 7.23: Adapted with permission from San Miguel, P., Tikhonov, A., Jin, Y. K., Motchoulskaia, N., Zakharov, D., Melake-Berhan, A., Springer, P. S., Edwards, K. J., Lee, M., Avramova, Z., et al. (1996) Nested retrotransposons in the intergenic regions of the maize genome. *Science* 274: 765–768. Copyright © 1996 American Association for the Advancement of Science.

Figure 7.25: O. L. Miller, B. R. Beatty, D. W. Fawcett/Visuals Unlimited, Swanzey, NH.

Figure 7.35: Figure adapted from McClintock, B. (1952) Chromosome organization and genic expression. *Cold Spring Harbor Symp. Quant. Biol.* 16: 13–47. Used with permission from Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. Legend from Keller, E. F. (1983) *A Feeling for the Organism: The Life and Work of Barbara McClintock*. W. H. Freeman, New York, p. 82. Copyright © 1983. Used with permission.

Figure 7.36: Photographs from F. W. Goro, published in Federov, N. V. (1984) Transposable genetic elements in maize. *Sci. Am.* 250(6): 84–98. Used with permission of the Estate of F. W. Goro.

Figure 7.37: Keller, J., Lim, E., James, D. W. Jr., Dooner, H. K. (1992) Germinal and somatic activity of the maize element activator (*Ac*) in *Arabidopsis*. *Genetics* 131: 449–459. Used with permission from the Genetic Society of America.

Figure 7.38: Neuffer, M. G., Coe, E. H., Wessler, S. R. (1997) *Mutants of Maize*. Cold

Spring Harbor Laboratory Press, Cold Spring Harbor, NY.

Figure 7.39: Chuck, G., Lincoln, C., Hake, S. (1996) *KNAT1* induces lobed leaves with ectopic meristems when overexpressed in *Arabidopsis*. *Plant Cell* 8: 1277–1289.

Figure 7.40: Taylor, J. E., Renwick, K. F., Webb, A. A., McAinsh, M. R., Furini, A., Bartels, C., Quatrano, R. S., Marcotte, W. R. Jr., Hetherington, A. M. (1995) ABA-regulated promoter activity in stomatal guard cells. *Plant J.* 7: 129–134. Used with permission from Blackwell Science Ltd.

Figure 7.47A: Reprinted by permission from Luger, K., Mäder, A. W., Richmond, R. K., Sargent, D. F., Richmond, T. J. (1997) Crystal structure of the nucleosome core particle at 2.8 Å resolution. *Nature* 389: 251–260. Copyright © 1997 Macmillan Magazines Ltd.

Figure 7.47B,C: Karp, G. (1996) *Cell and Molecular Biology: Concepts and Experiments*. John Wiley & Sons, New York. Used with permission from John Wiley & Sons. Copyright © 1996.

Figure 7.48: Paul, A.-L., Vasil, V., Vasil, I. K., Ferl, R. J. (1987) Constitutive and anaerobically induced DNase-I hypersensitive sites in the 5' region of the maize *ADHI* gene. *Proc. Natl. Acad. Sci. USA* 84: 799–803. Copyright © 1987 National Academy of Sciences, U.S.A.

Figure 7.50: Electronmicrograph from Paulson, J. R., Laemmli, U. K. (1977) The structure of histone-depleted metaphase chromosomes. *Cell* 12: 817–828. Copyright © 1977 Cell Press.

Figure 7.51: Kermicle, J. L. (1996) Epigenetic silencing and activation of a maize *r* gene. In *Cold Spring Harbor Monograph Series, 32. Epigenetic mechanisms of gene regulation*, V.E.A. Russo, R. A. Martienssen, and A. D. Riggs, eds. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, pp. 267–287.

Figure 7.52: Hollick, J. B., Dorweiler, J. E., Chandler, V. L. (1997) Paramutation and related allelic interactions. Reprinted from *Trends Genet.* 13: 302–308, Copyright © 1997 with permission from Elsevier Science.

Figure 7.53: Meyer, P. (1996) DNA methylation of flower color transgenes in *Petunia hybrida*. In *Cold Spring Harbor Monograph Series, 32. Epigenetic mechanisms of gene regulation*, V.E.A. Russo, R. A. Martienssen, and A. D. Riggs, eds. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, pp. 305–317.

Figure 7.54A: Jorgenson, R. (1993) The germinal inheritance of epigenetic information in plants. *Philos. Trans. R. Soc. Lond. B* 339: 173–181. Reprinted by permission of the President and Council of the Royal Society.

Figure 7.54B: Jorgenson, R. A., Napoli, C. A. (1996) A responsive regulatory system is revealed by sense suppression of pigment

genes in *Petunia* flowers. In *Genomes: Proceedings of the 22nd Stadler Genetics Symposium*, J. P. Gustafson and R. B. Flavell, eds. Plenum Press, New York, pp. 159–176.

Figure 7.54C: Napoli, C., Lemieux, C., Jorgenson, R. (1990) Introduction of a chimeric chalcone synthase gene into *petunia* results in reversible co-suppression of homologous genes in trans. *Plant Cell* 2: 279–289.

Figure 7.54D: Jorgenson, R. (1994) Developmental significance of epigenetic impositions on the plant genome: a paragenetic function for chromosomes. *Dev. Genet.* 15: 523–532. Used with permission from John Wiley & Sons, Inc. Copyright © 1994.

Figure 7.56: Reprinted with permission from Ronemus, M. J., Galbiati, M., Ticknor, C., Chen, J., Dellaporta, S. L. (1996) Demethylation-induced developmental pleiotropy in *Arabidopsis*. *Science* 273: 654–657. Copyright © 1996 American Association for the Advancement of Science.

Box 7.5 (left page): Paul, A.-L., Ferl, R. L. (1991) In vivo footprinting reveals unique cis-elements and different modes of hypoxic induction in maize *Adh1* and *Adh2*. *Plant Cell* 3: 159–168.

Box 7.5A,C: Paul, A.-L., Ferl, R. L. (1994) In vivo footprinting identifies an activating element of the maize *Adh2* promoter specific for root and vascular tissues. *Plant J.* 5: 523–533. Used with permission from Blackwell Science Ltd.

Chapter 8

Figure 8.3: Adapted from Lam, H.-M., Coschigano, K., Schultz, C., Melo-Oliveira, R., Tjaden, G., Oliveira, I., Ngai, N., Hsieh, M.-H., Coruzzi, G. (1995) Use of *Arabidopsis* mutants and genes to study amide amino acid biosynthesis. *Plant Cell* 7: 887–898.

Figure 8.8: Adapted from Wallgrove, R. M., Turner, J. C., Hall, N. P., Kendall, A. C., Bright, S.W.J. (1987) Barley mutants lacking chloroplast glutamine synthetase—biochemical and genetic analysis. *Plant Physiol.* 83: 155–158.

Figure 8.9: Edwards, J. W., Walker, E. L., Coruzzi, G. M. (1990) Cell-specific expression in transgenic plants reveals nonoverlapping roles for chloroplast and cytosolic glutamine synthetase. *Proc. Natl. Acad. Sci. USA* 87: 3459–3463. Used with permission.

Figure 8.11: Photographs: Coschigano, K. T., Melo-Oliveira, R., Lim, J., Coruzzi, G. M. (1998) *Arabidopsis gls* mutants and distinct Fd-GOGAT genes: implications for photorespiration and primary nitrogen assimilation. *Plant Cell* 10: 741–752.

Figure 8.12: Electrophoretogram: Coschigano, K. T., Melo-Oliveira, R., Lim, J., Coruzzi, G. M. (1998) *Arabidopsis gls* mutants and distinct Fd-GOGAT genes: implications for photorespiration and primary nitrogen assimilation. *Plant Cell* 10: 741–752.

Figure 8.15: Melo-Oliveira, R., Oliveira, I. C., Coruzzi, G. M. (1996) *Arabidopsis* mutant analysis and gene regulation define a nonredundant role for glutamate dehydrogenase in nitrogen assimilation. *Proc. Natl. Acad. Sci. USA* 93: 4718–4723. Copyright © 1996 National Academy of Sciences, U.S.A.

Figure 8.18A,B: Schultz, C. J., Hsu, M., Miesak, B., Coruzzi, G. M. (1998) *Arabidopsis* mutants define an in vivo role for isoenzymes of aspartate aminotransferase in plant nitrogen assimilation. *Genetics* 149: 491–499. Copyright © 1998 Genetic Society of America.

Figure 8.22: Eckes, P., Schmitt, P., Winfried, D., Wegenmeyer, F. (1989) Overproduction of alfalfa glutamine synthetase in transgenic tobacco plants. *Mol. Gen. Genet.* 217: 263–268. Copyright © 1989 Springer-Verlag.

Figure 8.23A,B: Adapted (A) and reprinted (B) by permission from Kozaki, A., Takeba, G. (1996) Photorespiration protects C₃ plants from photooxidation. *Nature* 384: 557–560. Copyright © 1996 Macmillan Magazines Ltd.

Figure 8.26: Adapted from Dyer, W. E., Weaver, L. M., Zhao, J., Kuhn, D. N., Weller, S. C., Herrmann, K. M. (1990) A complementary DNA encoding 3-deoxy-D-arabino-heptulosonate-7-phosphate synthase from *Solanum tuberosum* L. *J. Biol. Chem.* 265: 1608–1614. Copyright © 1990 American Society for Biochemistry and Molecular Biology.

Figure 8.37: R. Last, Cereon Genomics LLC, Cambridge, MA; previously unpublished.

Figure 8.41A: Hyde, C. C., Ahmed, S. A., Padlan, E. A., Miles, E. W., Davies, D. R. (1998) Three-dimensional structure of the tryptophan synthase $\alpha_2\beta_2$ multienzyme complex from *Salmonella typhimurium*. *J. Biol. Chem.* 263: 17857–17871. Copyright © 1988 American Society for Biochemistry and Molecular Biology.

Figure 8.41B: Adapted from Miles, E. W. (1995) Tryptophan synthase structure, function, and protein engineering. In *Subcellular Biochemistry: Proteins: Structure, Function, and Engineering*, B. B. Biswas and S. Roy, eds. Plenum Press, New York, pp. 207–254.

Figure 8.43A: Graph adapted from Dyer, W. E., Henstrand, J. M., Handa, A. K., Herrmann, K. M. (1989) Wounding induces the first enzyme of the shikimate pathway in Solanaceae. *Proc. Natl. Acad. Sci. USA* 86: 7370–7373. Used with permission.

Figure 8.43B: Immunoblot adapted from Zhao, J., Last, R. L. (1996) Coordinate regulation of the tryptophan biosynthetic pathway and idolic phytoalexin accumulation in *Arabidopsis*. *Plant Cell* 8: 2235–2244.

Figure 8.55: Biou, V., Dumas, R., Cohen-Addad, C., Douce, R., Job, D., Pebay-Peroula, E. (1997) The crystal structure of plant acetoxy acid isomer or reductase complexed with NADPH, two magnesium ions,

and a herbicidal transition site analog determined at 1.645 Å resolution. *EMBO J.* 16: 3405–3415. Used with permission from Oxford University Press.

Box 8.3: Dutch National Asparagus and Mushroom Museum, Horst-Melderslo, The Netherlands.

Box 8.4: Padgett, S. R., Re., D. B., Barry, G. F., Eicholtz, D. E., Delannay, X., Fuchs, R. C., Kishore, G. M., Fraley, R. T. (1996) New weed control opportunities: development of soybeans with a Roundup Ready gene. In *Herbicide-Resistant Crops: Agricultural, Environmental, Economic, Regulatory, and Technical Aspects*, S. O. Duke, ed. CRC Press, Boca Raton, FL, pp. 53–84. Copyright © 1996 CRC Press.

Box 8.5B: R. Last, Cereon Genomics LLC, Cambridge, MA, previously unpublished.

Box 8.6B: Bohlmann, J., De Luca, V., Eilert, U., Martin, W. (1995) Purification and cDNA cloning of anthranilate synthase from *Ruta graveolens*: modes of expression and properties of native and recombinant enzymes. *Plant J.* 7: 491–501. Used with permission from Blackwell Science Ltd.

Box 8.10A: Frankard, V., Ghisla, M., Jacobs, M. (1992) Two feedback-insensitive enzymes of the aspartate pathway in *Nicotiana sylvestris*. *Plant Physiol.* 99: 1285–1293.

Table 8.1: *Arabidopsis* data from Somerville, C. R., Ogren, W. L. (1980) Inhibition of photosynthesis in *Arabidopsis* mutants lacking leaf glutamate synthase activity. *Nature* 286: 257–259. Copyright © 1980 Macmillan Magazines Ltd. Barley data from Kendall, A. C., Wallsgrove, R. M., Hall, N. P., Turner, J. C., Lea, P. J. (1986) Carbon and nitrogen metabolism in barley (*Hordeum vulgare*) mutants lacking ferredoxin-dependent glutamate synthase. *Planta* 168: 316–323. Copyright © 1986 Springer-Verlag.

Chapter 9

Figure 9.6: Sussman, J. L., Holbrook, S. R., Warrant, R. W., Church, G. M., Kim, S.-H. (1978) Crystal structure of yeast phenylalanine t-RNA. I. Crystallographic refinement. *J. Mol. Biol.* 123: 607. Reprinted by permission of Academic Press. Copyright © 1978.

Figure 9.7: Reprinted by permission from Perona, J. J., Rould, M. A., Steitz, T. A. (1991) Structural basis of anticodon loop recognition by glutamyl transfer RNA synthetase. *Nature* 352: 213–218. Copyright © 1991 Macmillan Magazines Ltd.

Figure 9.9: Hill, W. E. (1990) Probing ribosome structure and function by using short complementary DNA oligomers. In *The Ribosome: Structure, Function, and Evolution*, W. E. Hill, J. Weller, T. Gluick, C. Merryman, R. T. Marconi, A. Tassanahajohn, and W. E. Teppich, eds. ASM Press, Washington, DC, pp. 253–262. Used with permission from the American Society for Microbiology.

Figure 9.32B: Baneyx, F., Bertsch, U., Kalbach, C. E., van Der Vies, S. M., Soll, J., Gatenby, A. A. (1995) Spinach chloroplast cpn 21 co-chaperonin possesses two functional domains fused together in a toroidal structure and exhibits nucleotide-dependent binding to plastid chaperonin 60. *J. Biol. Chem.* 270: 10695–10702. Copyright © 1995 American Society for Biochemistry and Molecular Biology.

Figure 9.33: Bukau, B., Horwich, A. L. (1998) The Hsp70 and Hsp60 chaperone machines. *Cell* 92: 351–366. Copyright © 1998 Cell Press.

Figure 9.38: I. Andersson, Swedish University of Agricultural Sciences, Uppsala; previously unpublished.

Figure 9.43B: Vierstra, R. D. (1996) Proteolysis in plants—mechanisms and functions. *Plant Mol. Biol.* 32: 275–302. Used with kind permission from Kluwer Academic Publishers.

Box 9.3: Mitsuko Williams, University of Illinois at Urbana-Champaign; previously unpublished.

Box 9.4: Electrophoretogram from Danon, A., Mayfield, S., Light regulated translational activators: identification of chloroplast gene-specific mRNA binding proteins. (1991) *EMBO J.* 10: 3993–4001. Used with permission from Oxford University Press.

Chapter 10

Figure 10.2: Adapted from Browse, J., Somerville, C. R. (1994) Glycerolipids. In *Arabidopsis*, E. Meyerowitz and C. R. Somerville, eds. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, pp. 881–912. Used with permission from Cold Spring Harbor Laboratory Press.

Figure 10.4: Reprinted with permission from Post-Beittenmiller, D. (1996) Biochemistry and molecular biology of wax production in plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 47: 405–430. Copyright © 1996 Annual Reviews. www.AnnualReviews.org

Figure 10.6: Reprinted with permission from Venable, R. M., Zhang, Y., Hardy, B. J., Pastor, R. W. (1993) Molecular dynamics simulation of a lipid bilayer and of hexadecane: an investigation of membrane fluidity. *Science* 262: 223–226. Copyright © 1993 American Association for the Advancement of Science.

Figure 10.17: Barton, P. G., Gunstone, F. D. (1975) Hydrocarbon chain packing and molecular motion in phospholipid bilayers formed from unsaturated lecithins. Synthesis and properties of 16 positional isomers of 1,2-dioctadecenoyl-sn-glycero-3-phosphorylcholine. *J. Biol. Chem.* 250: 4470–4476. Copyright © 1975 American Society for Biochemistry and Molecular Biology.

Figure 10.19A: J. Shanklin, Brookhaven National Laboratory, Upton, NY; previously unpublished.

Figure 10.19B,C: Lindqvist, Y., Huang, W., Schneider, G., Shanklin, J. (1996) Crystal structure of Δ -9 stearoyl-acyl carrier protein desaturase from castor seed and its relationship to other di-iron proteins. *EMBO J.* 15: 4081–4092. Used with permission from Oxford University Press.

Figure 10.21: Adapted from Browse, J., Somerville, C. R. (1994) Glycerolipids. In *Arabidopsis*, E. Meyerowitz and C. R. Somerville, eds. Cold Spring Harbor Laboratory Press, Cold Spring Harbor Laboratory Press, NY, pp. 881–912. Used with permission from Cold Spring Harbor Laboratory Press.

Figure 10.23: Vigh, L., Los, D. A., Horvath, I., Murata, N. (1993) The primary signal in the biological perception of temperature: Pd-catalyzed hydrogenation of membrane lipids stimulated the expression of the *desA* gene in *Synechocystis* PCC6803. *Proc. Natl. Acad. Sci. USA* 90: 9090–9094. Copyright © 1993 National Academy of Sciences, U.S.A.

Figure 10.28: Data from Sauer, F. D., Kramer, J.K.G. (1983) The problems associated with the feeding of high erucic acid rapeseed oils and some fish oils to experimental animals. In *High and Low Erucic Acid Rapeseed Oils: Production, Usage, Chemistry, and Toxicological Evaluation*, J.K.G. Kramer, F. D. Sauer, and W. J. Pidgen, eds. Academic Press, New York, pp. 253–292. Used with permission from Academic Press. Copyright © 1983.

Figure 10.31: Browse, J., Somerville, C. R. (1994) Glycerolipids. In *Arabidopsis*, E. Meyerowitz and C. R. Somerville, eds. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, pp. 881–912. Used with permission from Cold Spring Harbor Laboratory Press.

Figure 10.33: Kunst, L., Browse, J., Somerville, S. R. (1988) Altered regulation of lipid biosynthesis in a mutant of *Arabidopsis* deficient in chloroplast glycerol phosphate acyltransferase activity. *Proc. Natl. Acad. Sci. USA* 85: 4143–4147. Used with permission.

Figure 10.42: C. Somerville, Carnegie Institute of Washington, Stanford, CA; previously unpublished.

Figure 10.43A–E: Lightner, J., Sames, D. W. Jr., Donner, H. K., Browse, J. (1994) Altered body morphology is caused by increased stearate levels in a mutant of *Arabidopsis*. *Plant J.* 6: 401–412. Reprinted with permission from Blackwell Science Ltd.

Figure 10.44A,B: McConn, M., Browse, J. (1996) The critical requirement for linolenic acid is pollen development, not photosynthesis, in an *Arabidopsis* mutant. *Plant Cell* 8: 403–416.

Figure 10.44C: McConn, M., Browse, J. (1998) Polyunsaturated membranes are required for photosynthetic competence in a mutant of *Arabidopsis*. *Plant J.* 15: 521–530. Reprinted with permission from Blackwell Science Ltd.

Figure 10.45: C. Somerville, Carnegie Institute of Washington, Stanford, CA; previously unpublished results.

Figure 10.46: Adapted from Janiak, M. J., Small, D. M., Shipley, G. G. (1976) Nature of the thermal pretransition of synthetic phospholipids: dimyristoyl- and dipalmitoyllecithin. *Biochemistry* 15: 4574–4580. Copyright © 1976 American Chemical Society.

Figure 10.47A: Hugly, S., Somerville, C. R. (1992) A role for membrane lipid polyunsaturation in biogenesis at low temperature. *Plant Physiol.* 99: 197–202.

Figure 10.47B: Miquel, M., James, D. Jr., Dooner, H., Browse, J. (1993) *Arabidopsis* requires polyunsaturated lipids for low temperature survival. *Proc. Natl. Acad. Sci. USA* 90: 6208–6212. Copyright © 1993 National Academy of Sciences, U.S.A.

Figure 10.47C,D: Wu, J., Lightner, J., Warwick, N., Browse, J. (1997) Low-temperature damage and subsequent recovery of *fab1* mutant *Arabidopsis* exposed to 2°C. *Plant Physiol.* 113: 347–356.

Figure 10.48: Mike Thomashow, Michigan State University, East Lansing; previously unpublished.

Figure 10.53: McConn, M., Creelman, R. A., Bell, E., Mullet, J. E., Browse, J. (1997) Jasmonate is essential for insect defense in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* 94: 5473–5477. Copyright © 1997 National Academy of Sciences, U.S.A.

Figure 10.54A: Riederer, M., Schonherr, J. (1988) Development of plant cuticles: fine structure and cutin composition of *Clivia miniata* Reg. leaves. *Planta* 174: 127–138. Copyright © 1988 Springer-Verlag.

Figure 10.54B: Lenzian, K. J., Schonherr, J. (1983) In vivo study of cutin synthesis in leaves of *Clivia miniata* Reg. *Planta* 158: 70–75. Copyright © 1983 Springer-Verlag.

Figure 10.55: Jeffree, C. E. (1996) Structure and ontogeny of plant cuticles. In *Plant Cuticles: An Integrated Functional Approach*, G. Kerstiens, ed. BIOS Scientific Publishers, Oxford, UK, pp. 33–82. Used with permission of BIOS Scientific Publishers Ltd.

Figure 10.56: Riederer, M., Schonherr, J. (1988) Development of plant cuticles: fine structure and cutin composition of *Clivia miniata* Reg. leaves. *Planta* 174: 127–138. Copyright © 1988 Springer-Verlag.

Figure 10.59: Sutter, E. (1984) Chemical composition of epicuticular wax in cabbage plants grown in vitro. *Can. J. Bot.* 62: 74–77. Reprinted with permission from NRC Research Press.

Figure 10.60: Jenks, M. A., Rich, P. J., Ashworth, E. N. (1994) Involvement of cork cells in the secretion of epicuticular wax filaments on *Sorghum bicolor* (L.) Moench. *Int. J. Plant Sci.* 155: 506–518. Reprinted with permission from the University of Chicago Press.

- Figure 10.62: D. Preuss, University of Chicago, Illinois; previously unpublished.
- Figure 10.63: Huang, A.H.C. (1987) Lipases. In *The Biochemistry of Plants: A Comprehensive Treatise. Vol. 9, Lipids: Structure and Function*, P. K. Stumpf, ed. Academic Press, New York, pp. 91–119. Reprinted by permission of Academic Press. Copyright © 1987.
- Figure 10.65A: Fernandez, D. E., Staehelin, L. A. (1987) Does gibberellic acid induce the transfer of lipase from protein bodies to lipid bodies in barley aleurone cells? *Plant Physiol.* 85: 487–496.
- Figure 10.65B: Reprinted with permission from Huang, A.H.C. (1992) Oil bodies and oleosins in seeds. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 43: 177–200. Copyright © 1992 Annual Reviews. www.annualreviews.org
- Figure 10.65C: Qu, R., Huang, A.H.C. (1990) Oleosin kD18 on the surface of oil bodies in maize: genomic and complementary DNA sequences and the deduced protein structure. *J. Biol. Chem.* 6: 2238–2243. Copyright © 1990 American Society for Biochemistry and Molecular Biology.
- Figure 10.66: Adapted with permission from Huang, A.H.C. (1992) Oil bodies and oleosins in seeds. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 43: 177–200. Copyright © 1992 Annual Reviews. www.annualreviews.org
- Figure 10.69: Ackman, R. G. (1983) Chemical composition of rapeseed oil. In *High and Low Erucic Acid Rapeseed Oils*, J.K.G. Kramer, F. D. Sauer, and W. J. Pidgen, eds. Academic Press, Toronto, pp. 85–129. Used with permission from Academic Press. Copyright © 1983.
- Figure 10.69 Inset: Illustration by B. E. Nicholson. Reprinted from *The Oxford Book of Food Plants*. (1969) by permission of Oxford University Press.
- Figure 10.70: Based on data from Roesler, K. R., Shintani, D., Savage, L., Boddupalli, S., Ohlrogge, J. B. (1997) Targeting of the *Arabidopsis* homomeric acetyl-CoA carboxylase to *Brassica napus* seed plastids. *Plant Physiol.* 13: 75–81.
- Figure 10.71: J. Ohlrogge, Michigan State University, East Lansing; previously unpublished.
- Figure 10.73A: B. Page, University of Alberta, Edmonton, Canada; previously unpublished.
- Figure 10.75: Nawrath, C., Poirier, Y., Somerville, C. R. (1994) Targeting of the polyhydroxybutyrate biosynthetic pathway to the plastids of *Arabidopsis thaliana* results in high levels of polymer accumulation. *Proc. Natl. Acad. Sci. USA* 91: 12760–12764. Copyright © 1994 National Academy of Sciences, U.S.A.
- Box 10.1: J. Ohlrogge, Michigan State University, East Lansing; previously unpublished.
- Box 10.4: C. Somerville, Carnegie Institute of Washington, Stanford, CA; previously unpublished.
- Table 10.5: Data from Battey, J. F., Schmid, K. M., Ohlrogge, J. B. (1989) Genetic engineering for plant oils: potential and limitations. *Trends Biotechnol.* 7: 122–126. Copyright © 1989 Elsevier Science.
- ## Chapter 11
- Chapter 11 opening photograph: P. Heslop-Harrison, John Innes Centre, Norwich, UK; previously unpublished.
- Figure 11.1: Photographs: Russel, P., Nurse, P. (1987) Negative regulation of mitosis by *Wee1*, a gene encoding a protein kinase homolog. *Cell* 49: 559–567. Copyright © 1987 Cell Press.
- Figure 11.5B: Minshull, J. (1989) The role of cyclin synthesis modification and destruction in the control of cell division. In *The Cell Cycle*, R. Brooks et al., eds. *J. Cell Sci. Suppl.* 12: 77–98. Used with permission from Company of Biologists Ltd.
- Figure 11.6B: Szabados, I., Dudits, D. (1980) Fusion between interphase and mitotic plant protoplasts. Induction of premature chromosome condensation. *Exp. Cell Res.* 127: 441–446.
- Figure 11.9: Adapted from Reichheld, J. P., Gigot, C., Chaubet-Gigot, N. (1998) Multi-level regulation of histone gene expression during the cell cycle in tobacco cells. *Nucleic Acids Res.* 26: 3255–3262. Used by permission of Oxford University Press.
- Figure 11.10: Photographs: Sitte, P., Ziegler, H., Ehrendorfer, F., Bresinsky, A., eds. (1998) *Strasburger Lehrbuch der Botanik*, 34th ed. Gustav Fischer Verlag. Copyright © 1998 Spektrum Akademischer Verlag, Heidelberg.
- Figure 11.11: Ribbon structures: CDK—Reprinted by permission from de Bondt, H. L., Rosenblatt, J., Jancarik, J., Jones, H. D., Morgan, D. O., Kim, S.-H. (1993) Crystal structure of cyclin-dependent kinase 2. *Nature* 363: 595–602. Copyright © 1993 Macmillan Magazines Ltd. CDK2–cyclin A complex—Reprinted by permission from Jeffrey, P. D., Russo, A. A., Polyak, K., Gibbs, E., Hurwitz, J., Massague, J., Pavletich, N. P. (1995) Mechanism of CDK activation revealed by the structure of a cyclinA–CDK2 complex. *Nature* 376: 313–320. Copyright © 1995 Macmillan Magazines Ltd. T160-phosphorylated CDK2–cyclinA complex—Reprinted by permission from Russo, A. A., Jeffrey, P. D., Pavletich, N. P. (1996) Structural basis of cyclin-dependent kinase activation by phosphorylation. *Nat. Struct. Biol.* 3: 696–700. Copyright © 1996 Nature Publishing Group, New York. CDK2–cycA–CKI complex—Reprinted by permission from Russo, A. A., Jeffrey, P. D., Patten, A. K., Massague, J., Pavletich, N. P. (1996) Crystal structure of the p27^{Kip1} cyclin-dependent kinase inhibitor bound to the cyclin A–Cdk2 complex. *Nature* 382: 325–331. Copyright © 1996 Macmillan Magazines Ltd.
- Figure 11.18B,C: Ferreria, P.C.G., Hemerly, A. S., Engler, J.D.A., Van Montagu, M., Inzè, D., Engler, G. (1994) Developmental expression of the *Arabidopsis* cyclin gene *cyc1AT*. *Plant Cell* 6: 1763–1774.
- Figure 11.19A: Lyndon, R. F. (1998) *The Shoot Apical Meristem: Its Growth and Development*. Cambridge University Press, Cambridge, UK. Reprinted with permission from the Cambridge University Press.
- Figure 11.20A: Colón-Carmona, A., You, R., Haimovich-Gal, T., Doerner, P. (1999) Spatio-temporal analysis of mitotic activity with a labile cyclin–GUS fusion protein. *Plant J.* 20: 503–508. Used with permission from Blackwell Science Ltd.
- Figure 11.20B: Clowes, originally published in O'Brien, T. P., McCully, M. E. (1969) *Plant Structure and Development—A Pictorial and Physiological Approach*. Collier-Macmillan, Ltd. London.
- Figure 11.22A: Reprinted from Yokota, T. (1997) The structure, biosynthesis and function of brassinosteroids. *Trends Plant Sci.* 2: 137–143. Copyright © 1997 with permission from Elsevier Science.
- Figure 11.22B: Boerjan, W., Cervera, M.-T., Delarue, M., Beeckman, T., Van Montagu, M., Inzè, D. (1995) *superrroot*, a recessive mutation in *Arabidopsis*, confers auxin overproduction. *Plant Cell* 7: 1405–1419.
- Figure 11.22C: McCarty, D. R., Hattori, T., Carson, C. B., Vasil, V., Lazar, M., Vasil, I. K. (1991) The *Viviparous-1* developmental gene of maize encodes a novel transcription activator. *Cell* 66: 895–905. Copyright © 1991 Cell Press.
- Figure 11.24: Top: Reprinted by permission from Doerner, P. D., Jorgensen, J.-E., You, R., Steppuhn, J., Lamb, C. (1996) Control of root growth and development by cyclin expression. *Nature* 380: 520–523. Copyright © 1996 Macmillan Magazines Ltd. Bottom: P. Doerner, University of Edinburgh, Scotland; previously unpublished.
- Figure 11.25A: Reprinted by permission from Doerner, P. D., Jorgensen, J.-E., You, R., Steppuhn, J., Lamb, C. (1996) Control of root growth and development by cyclin expression. *Nature* 380: 520–523. Copyright © 1996 Macmillan Magazines Ltd.
- Figure 11.25B: Hemerly, A., Engler, J.D.A., Bergounioux, C., Van Montagu, M., Engler, G., Inzè, D., Ferreira, P. (1995) Dominant negative mutants of the Cdc2 kinase uncouple cell division from iterative plant development. *EMBO J.* 14: 3925–3936.
- Figure 11.26A: S. Smith, University of Edinburgh, Scotland; previously unpublished.
- Figure 11.26B: Raven, P. H., Evert, R. F., Eichhorn, S. E., eds. (1992) *Biology of Plants*, 5th ed. Worth Publishers, New York. Used with permission.
- Figure 11.27: Adapted from Galbraith, D. W., Harkins, K. R., Knapp, S. (1991) Systemic

endopolyploidy in *Arabidopsis thaliana*. *Plant Physiol.* 96: 985–989.

Box 11.3: Reprinted with permission from Koshland, D.E.J., Goldbeter, S., Stock, J. B. (1984) Amplification and adaptation in regulatory and sensory systems. *Science* 217: 220–225. Copyright © 1984 American Association for the Advancement of Science.

Part 3 opening photograph:

Galen Rowell/Mountain Light Photography, Emeryville, CA.

Chapter 12

Figure 12.1B: Staehelin, L. A., van der Staay, G.W.M. (1996) Structure, composition, functional organization and dynamic properties of thylakoid membranes. In *Oxygenic Photosynthesis: The Light Reactions*, D. R. Ort and C. F. Yocum, eds. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp. 11–30. Copyright © 1996. Used with kind permission of Kluwer Academic Publishers.

Figure 12.8: Cunningham, F. X. Jr., Gantt, E. (1998) Genes and enzymes of carotenoid biosynthesis in plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 49: 557–583. Copyright © 1998 Annual Reviews. www. AnnualReviews.org

Figure 12.10A,B: Lancaster, C.P.D., Michel, H. (1996) Three-dimensional structures of photosynthetic reaction centers. *Photosynth. Res.* 48: 65–740. Copyright © 1996. Used with kind permission of Kluwer Academic Publishers.

Figure 12.13: Adapted from He, W. Z. (1998) Photosystems I and II. In *Photosynthesis: A Comprehensive Treatise*, A. S. Reghavra, ed. Cambridge University Press, New York and Cambridge, pp. 29–43. Reprinted with the permission of Cambridge University Press.

Figure 12.14: Schubert, W. D., Klukas, O., Krauss, N., Saenger, W., Fromme, P., Witt, H. T. (1997) Photosystem I of *Synechococcus elongatus* at 4 Å resolution: comprehensive structure analysis. *J. Mol. Biol.* 272: 741–769. Reprinted by permission of Academic Press. Copyright © 1997.

Figure 12.15: Schubert, W. D., Klukas, O., Krauss, N., Saenger, W., Fromme, P., Witt, H. T. (1997) Photosystem I of *Synechococcus elongatus* at 4 Å resolution: comprehensive structure analysis. *J. Mol. Biol.* 272: 741–769. Reprinted by permission of Academic Press. Copyright © 1997.

Figure 12.16A: Reprinted by permission from Kühlbrandt, W., Wang, D. N., Fujiyoshi, Y. (1994) Atomic model of plant light harvesting complex by electron crystallography. *Nature* 367: 614–621. Copyright © 1994 Macmillan Magazines Ltd.

Figure 12.16B: Reprinted from Kühlbrandt, W. (1994) Structure and function of the plant light harvesting complex, LHC-II. *Curr. Opin. Struct. Biol.* 4: 519–528. Copyright © 1994 with permission from Elsevier Science.

Figure 12.17: Adapted from Jansson, S. (1994) The light harvesting chlorophyll *a/b* binding proteins. *Biochim. Biophys. Acta* 1184: 1–19. Copyright © 1994 with permission from Elsevier Science.

Figure 12.23: Adapted from He, W. Z. (1998) Photosystems I and II. In *Photosynthesis: A Comprehensive Treatise*, A. S. Reghavra, ed. Cambridge University Press, New York and Cambridge, pp. 29–43. Reprinted with the permission of Cambridge University Press.

Figure 12.26: J. Fernandez-Velasco, University of California, Berkeley, and E. Berry, Ernest O. Lawrence Berkeley National Laboratory; previously unpublished.

Figure 12.27: Carrell, C. J., Zhang, H., Cramer, W. A., Smith, J. L. (1997) Biological identity and diversity in photosynthesis and respiration: structure of the lumen-side domain of the chloroplast Rieske protein. *Structure* 5: 1613–1625. Used with permission from Current Biology Ltd.

Figure 12.28: Adapted from He, W. Z. (1998) Photosystems I and II. In *Photosynthesis: A Comprehensive Treatise*, A. S. Reghavra, ed. Cambridge University Press, New York and Cambridge, pp. 29–43. Reprinted with the permission of Cambridge University Press.

Figure 12.29: Babcock, G. T. (1973) Kinetics and intermediates in photosynthetic oxygen evolution. Ph.D. thesis, University of California, Berkeley.

Figure 12.31: Reprinted with permission from Yachandra, V. K., DeRose, V. J., Latimer, M. J., Mukerji, I., Sauer, K., Klein, M. P. (1993) Where plants make oxygen: a structural model for the photosynthetic oxygen-evolving manganese cluster. *Science* 260: 675–679. Copyright © 1993 American Association for the Advancement of Science.

Figure 12.35: Reprinted from Junge, W., Lill, H., Engelbrecht, S. (1997) ATP synthase: an electrochemical transducer with rotatory mechanics. *Trends Biochem. Sci.* 22: 420–423. Copyright © 1997 with permission from Elsevier Science.

Figure 12.36: Adapted from Cross, R. L., Duncan, T. M. (1996) Subunit rotator in F_0F_1 -ATP synthases as a means of coupling proton transport through F_0 to the binding changes in F_1 . *J. Bioenerg. Biomembr.* 28: 403–408. Copyright © 1996. Used with kind permission of Kluwer Academic Publishers.

Figure 12.37: Ernest O. Lawrence Berkeley National Laboratory, Photography and Digital Imaging Services; previously unpublished.

Figure 12.39: I. Andersson, Swedish University of Agricultural Sciences, Uppsala; previously unpublished.

Figure 12.45: E. H. Newcomb, University of Wisconsin, Madison; previously unpublished.

Figure 12.47: E. H. Newcomb, University of Wisconsin, Madison; previously unpublished.

Figure 12.48: Taiz, L., Zeiger, E. (1991) CO_2 Concentrating mechanisms II: The C_4 photosynthetic carbon assimilation (PCA) cycle. In *Plant Physiology* Benjamin/Cummings, Redwood City, CA. Copyright © 1991 The Benjamin/Cummings Publishing Co., Inc. Reprinted by permission of Addison Wesley Longman.

Box 12.2A: Melis, A. (1990) Regulation of photosystem stoichiometry in oxygenic photosynthesis. *Bot. Mag. Tokyo* 2 (Special Issue): 9–28. Copyright © 1990 Botanical Society of Japan.

Box 12.2B: R. Malkin, University of California, Berkeley; previously unpublished.

Box 12.3: The molecular models are from S. Dai and P. Schürmann, Université de Neuchâtel, Neuchâtel, Switzerland; previously unpublished.

Chapter 13

Chapter 13 opening photograph: Agricultural Research Service, US Department of Agriculture.

Box 13.3: Cover photograph for *Cell* 60 (1), by Bhattacharyya. Copyright © 1990 Cell Press.

Chapter 14

Figure 14.1B: R. Wiser, University of Wisconsin, Oshkosh; previously unpublished.

Figure 14.22B: Reprinted with permission from Iwata, S., Lee, J. W., Okada, K., Lee, J. K., Iwata, M., Rasmussen, B., Link, T. A., Ramaswamy, S., Jap, B. K. (1998) Complete structure of the 11-subunit bovine mitochondrial cytochrome bc_1 complex. *Science* 281: 64–71. Copyright © 1998 American Association for the Advancement of Science.

Figure 14.23B: Reprinted with permission from Tsukihara, T., Aoyama, H., Yamashita, E., Tomizaki, T., Yamaguchi, H., Shinzawa-Itoh, K., Nakashima, R., Yaono, R., Yoshikawa, S. (1996) The whole structure of the 13-subunit oxidized cytochrome-*c* oxidase at 2.8 Å. *Science* 272: 1136–1144. Copyright © 1996 American Association for the Advancement of Science.

Figure 14.44: Berry, J. A., Downton, J. S. (1982) Environmental regulation of photosynthesis. In *Photosynthesis, Development, Carbon Metabolism and Plant Productivity*, Vol. 2, Govindjee, ed. Academic Press, New York, pp. 263–343. Reprinted by permission of Academic Press. Copyright © 1982.

Figure 14.49: Berry, J. A., Downton, J. S. (1982) Environmental regulation of photosynthesis. In *Photosynthesis, Development, Carbon Metabolism and Plant Productivity*, Vol. 2, Govindjee, ed. Academic Press, New York, pp. 263–343. Reprinted by permission of Academic Press. Copyright © 1982.

Figure 14.50: Berry, J., Bjorkman, O. (1980) Photosynthetic response and adaptation to temperature in higher plants. *Annu. Rev. Plant Physiol.* 31: 491–543.

Box 14.2: M. Benstead, Fife, Scotland, UK; previously unpublished.

Part 4 opening photograph:

Galen Rowell/Mountain Light Photography, Emeryville, CA.

Chapter 15

Figure 15.1A–J, photograph: D. B. Fisher, Washington State University, Pullman; previously unpublished.

Figure 15.7: Protein data adapted from le Maire, M., Aggerbeck, L. P., Monteilhet, C., Andersen, J. P., Moller, J. V. (1986) The use of high-performance liquid chromatography for the determination of size and molecular weight of proteins: a caution and a list of membrane proteins suitable as standards. *Anal. Biochem.* 154: 525–535. Copyright © Academic Press. Dextran data adapted from Jorgensen, D. E., Moller, J. V. (1979) Use of flexible polymers as probes of glomerular pore size. *Am. J. Physiol.* 236: F103–F111. Copyright © American Physiological Society.

Figure 15.10A–D: Koontz, H., Biddulph, O. (1957) Factors regulating absorption and translocation of foliar applied phosphorus. *Plant Physiol.* 32: 463–470.

Figure 15.13A,B: Gunning, B.E.S., Pate, J. S., Green, L. W. (1970) Transfer cells in the vascular system of stems: taxonomy, associations with nodes, and structure. *Protoplasma* 71: 147–171. Copyright © 1970 Springer-Verlag.

Figure 15.16A,B: Sperry, J. S., Donnelly, J. R., Tyree, M. T. (1988) A method for measuring hydraulic conductivity and embolism in xylem. *Plant Cell Environ.* 11: 35–40. Used with permission from Blackwell Science Ltd.

Figure 15.18: Adapted by permission from Sperry, J. S. (1995) Limitations on stem water transport and their consequences. In *Plant Stems. Physiological and Functional Morphology*, B. L. Gartner, ed. Academic Press, San Diego, pp. 105–125. Copyright © 1995 Academic Press.

Figure 15.19: D. B. Fisher, Washington State University, Pullman; previously unpublished.

Figure 15.20A: Longitudinal section: D. B. Fisher, Washington State University, Pullman; previously unpublished. Cross-section: Overall, R. L., Wolfe, J., Gunning, B.E.S. (1982) Intercellular communication in *Azolla* roots: I. Ultrastructure of plasmodesmata. *Protoplasma* 111: 134–150. Copyright © 1982 Springer-Verlag.

Figure 15.21: Radford, J. E., Vesik, M., Overall, R. L. (1998) Callose deposition in plasmodesmata. *Protoplasma* 201: 30–37. Copyright © 1998 Springer-Verlag.

Figure 15.22: Radford, J. E., Vesik, M., Overall, R. L. (1998) Callose deposition in plasmodesmata. *Protoplasma* 201: 30–37. Copyright © 1998 Springer-Verlag.

Figure 15.23: Hepler, P. K. (1982) Endoplasmic reticulum in the formation of the cell plate and plasmodesmata. *Protoplasma* 111: 121–133. Copyright © 1982 Springer-Verlag.

Figure 15.25: Adapted from van Bell, A.J.E., Oparka, K. J. (1995) On the validity of plasmodesmograms. *Bot. Acta* 108: 174–182. Copyright © 1995 Georg Thieme Verlag, Stuttgart–New York.

Figure 15.26A–F: Adapted from Kollmann, R., Glockmann, C. (1991) Studies on graft unions. III. On the mechanism of secondary formation of plasmodesmata at the graft interface. *Protoplasma* 165: 71–85. Copyright © 1991 Springer-Verlag.

Figure 15.29: Adapted from Overall, R. L., Gunning, B.E.S. (1982) Intercellular communication in *Azolla* roots: II. Electrical coupling. *Protoplasma* 111: 151–160. Copyright © 1982 Springer-Verlag.

Figure 15.30A–D: Terry, B. R., Robards, A. W. (1987) Hydrodynamic radius alone governs the mobility of molecules through plasmodesmata. *Planta* 171: 145–157. Copyright © 1987 Springer-Verlag.

Figure 15.31: Duckett, C. M., Oparka, K. J., Prior, D.A.M., Dolan, L., Roberts, K. (1994) Dye-coupling in the root epidermis of *Arabidopsis* is progressively reduced during development. *Development* 120: 3247–3255. Used with permission from Company of Biologists Ltd.

Figure 15.32A,B: Wright, K. M., Oparka, K. J. (1997) Metabolic inhibitors induce symplastic movement of solutes from the transport phloem of *Arabidopsis* roots. *J. Exp. Bot.* 48: 1807–1814. Used with permission from Oxford University Press.

Figure 15.33A,B: Adapted from data of Tyree, M. T., Tammes, P.M.L. (1972) Translocation of uranin in the symplast of staminal hairs of *Tradescantia*. *Can. J. Bot.* 53: 2038–2046. Reprinted with permission from NRC Research Press.

Figure 15.34A,B: Yahalom, A., Warmbrodt, R. D., Laird, D. W., Traub, O., Revel, J. P., Willecke, K., Epel, B. L. (1991) Maize mesocotyl plasmodesmata proteins cross-react with connexin gap junction protein antibodies. *Plant Cell* 3: 407–417.

Figure 15.35A–C: Blackman, L. M., Boevink, P., Santa Cruz, S., Palukaitis, P., Oparka, K. J. (1998) The movement protein of cucumber mosaic virus traffics into sieve elements in minor veins of *Nicotiana glauca*. *Plant Cell* 10: 525–537.

Figure 15.36A,B: Fisher, D. B. (1975) Structure of functional soybean sieve elements. *Plant Physiol.* 56: 555–569.

Figure 15.37: Knoblauch, M., van Bel, A.J.E. (1998) Sieve tubes in action. *Plant Cell* 10: 35–50.

Figure 15.38: Adapted with permission from Watson, B. T. (1975) The influence of low temperature on the rate of translocation in the phloem of *Salix viminalis* L. *Ann. Bot.* 39: 889–900. Copyright © 1975 Academic Press.

Figure 15.39: Adapted with permission from Hocking, P. J. (1980) The composition of phloem exudate and xylem sap from tree tobacco (*Nicotiana glauca* Grah.). *Ann. Bot.* 45: 633–643. Copyright © 1980 Academic Press.

Figure 15.40: Esau, K., Thorsch, J. (1985) Sieve plate pores and plasmodesmata, the communication channels of the symplast: ultrastructural aspects and developmental relations. *Am. J. Bot.* 72: 1641–1653. Used with permission from Botanical Society of America, Inc. Copyright © 1985.

Figure 15.41 (left): Turgeon, R., Beebe, D. U., Gowan, E. (1993) The intermediary cell: minor-vein anatomy and raffinose oligosaccharide synthesis in the Scrophulariaceae. *Planta* 191: 446–456. Copyright © 1993 Springer-Verlag.

Figure 15.41 (right): Fisher, D. B., Housley, T. L., Christy, A. L. (1978) Source pool kinetics for ¹⁴C-photosynthate translocation in morning glory and soybean. *Plant Physiol.* 61: 291–295.

Figure 15.42A,B: Geiger, D. R., Sovonick, S. A., Shock, T. L., Fellows, R. J. (1974) Role of free space in translocation in sugar beet. *Plant Physiol.* 54: 892–898.

Figure 15.43A,B: Giaquinta, R. (1976) Evidence for phloem unloading from the apoplast. Chemical modification of membrane sulfhydryl groups. *Plant Physiol.* 57: 872–875.

Figure 15.44: Adapted from Wright, J. P., Fisher, D. B. (1981) Measurement of the sieve tube membrane potential. *Plant Physiol.* 67: 845–848.

Figure 15.45A,B: Turgeon, R., Wimmers, L. E. (1988) Different patterns of vein loading of exogenous [¹⁴C]sucrose in leaves of *Pisum sativum* and *Coleus blumei*. *Plant Physiol.* 87: 179–182.

Figure 15.46: Adapted from Russin, W. A., Evert, R. F. (1985) Studies on the leaf of *Populus deltoides* (Salicaceae): ultrastructure, plasmodesmatal frequency, and solute concentrations. *Am. J. Bot.* 72: 1232–1247. Used with permission from Botanical Society of America, Inc. Copyright © 1985.

Figure 15.48: Adapted from Schmalstig, J., Cosgrove, D. (1990) Coupling of solute transport and cell expansion in pea stems. *Plant Physiol.* 94: 1625–1633.

Figure 15.49: Oparka, K. J., Duckett, C. M., Prior, D.A.M., Fisher, D. B. (1994) Real-time imaging of phloem unloading in the root tip of *Arabidopsis*. *Plant J.* 6: 759–766. Reprinted with permission from Blackwell Science Ltd.

- Figure 15.51A–C: D. B. Fisher, Washington State University, Pullman; previously unpublished.
- Figure 15.52A,B: Fisher, D. B., Wang, N. (1993) A kinetic and microautoradiographic analysis of [¹⁴C]sucrose import by developing wheat grains. *Plant Physiol.* 101: 391–398.
- Figure 15.54A: Johnson, R. W., Dixon, M. A., Lee, D. R. (1992) Water relations of the tomato during fruit growth. *Plant Cell Environ.* 15: 947–953. Reprinted with permission from Blackwell Science Ltd.
- Figure 15.54B: Gandar, P. W., Turner, C. B. (1976) Potato leaf and tuber water potential measurements with a pressure chamber. *Am. Potato J.* 53: 1–14. Used with permission from the Potato Association of America.
- Figure 15.55: Wang, N., Fisher, D. B. (1994) The use of fluorescent tracers to characterize the post-phloem transport pathway in maternal tissues of developing wheat grains. *Plant Physiol.* 104: 17–27.
- Figure 15.56: Lang, A. (1990) Xylem, phloem and transpiration flows in developing apple fruits. *J. Exp. Bot.* 41: 645–651. Used with permission from Oxford University Press.
- Figure 15.57: Pritchard, J., Tomos, A. D. (1993) Correlating biophysical and biochemical control of root cell expansion. In *Water Deficits: Plant Responses from Cell to Community*, J.A.C. Smith and H. Griffiths, eds. BIOS Scientific Publishers, Oxford, UK, pp. 53–72. Used with permission of BIOS Scientific Publishers Ltd.
- Figure 15.58: Fisher, D. B., Wu, Y., Ku, M.S.B. (1992) Turnover of soluble proteins in the wheat sieve tube. *Plant Physiol.* 100: 1433–1441.
- Figure 15.59A,B: Oparka, K. J., Roberts, A. G., Boevink, P., Santa Cruz, S., Roberts, I., Pradel, K. S., Imlau, A., Kotlizky, G., Saure, N., Epel, B. (1999) Simple, but not branched, plasmodesmata allow the non-specific trafficking of proteins in developing tobacco leaves. *Cell* 97: 743–754. Copyright © 1999 Cell Press.
- Figure 15.59C,D: Imlau, A., Truernit, E., Sauer, N. (1999) Cell-to-cell and long-distance trafficking of the green fluorescent protein in the phloem and symplastic unloading of the protein into sink tissues. *Plant Cell* 11: 309–322.
- Figure 15.60: Fisher, D. B., Wu, Y., Ku, M.S.B. (1992) Turnover of soluble proteins in the wheat sieve tube. *Plant Physiol.* 100: 1433–1441.
- Figure 15.61A,B: Reprinted with permission from Xoconostle-Cazares, B., Xiang, Y., Ruiz-Medrano, R., Wang, H.-L., Monzer, J., Yoo, B. C., McFarland, K. C., Francheschi, V. R., Lucas, W. J. (1999) Plant paralog to viral movement protein that potentiates transport of mRNA into the phloem. *Science* 283: 94–98. Copyright © 1999 American Association for the Advancement of Science.
- Figure 15.63A,B: Reprinted with permission from Lucas, W. J., Bouche-Pillon, S., Jackson, D. P., Nguyen, L., Baker, L., Ding, B., Hake, S. (1995) Selective trafficking of KNOTTED-1 homeodomain protein and its mRNA through plasmodesmata. *Science* 270: 1980–1983. Copyright © 1995 American Association for the Advancement of Science.
- Figure 15.64: Ding, B. (1998) Intercellular protein trafficking through plasmodesmata. *Plant Mol. Biol.* 38: 279–310. With kind permission from Kluwer Academic Publishers.
- Figure 15.65: Adapted from Kragler, F., Monzer, J., Shash, K., Xoconostle-Cazares, B., Lucas, W. J. (1998) Cell-to-cell transport of proteins: requirement for unfolding and characterization of binding to a putative plasmodesmal receptor. *Plant J.* 15: 367–381. Reprinted with permission from Blackwell Science Ltd.
- Figure 15.66: Ding, B., Kwon, M.-O., Hammond, R., Owens, R. (1997) Cell-to-cell movement of potato spindle tuber viroid. *Plant J.* 12: 931–936. Reprinted with permission from Blackwell Science Ltd.
- Box 15.2C–E: D. B. Fisher, Washington State University, Pullman; previously unpublished.
- Box 15.3: Graph data from Holbrook, N. M. (1995) Negative xylem pressures in plants: a test of the balancing pressure technique. *Science* 270: 1193–1194. Copyright © 1995 American Association for the Advancement of Science.
- Box 15.4A: Eleftheriou, E. P., Tsekos, I. (1982) Development of protophloem in roots of *Aegilops comosa* var. *thessalica*. II. Sieve-element differentiation. *Protoplasma* 113: 221–233. Copyright © 1982 Springer-Verlag.
- Box 15.4B,C: Eleftheriou, E. P. (1993) Differentiation of abnormal sieve elements in roots of wheat (*Triticum aestivum* L.) affected by colchicine. *New Phytol.* 125: 813–827. Reprinted with permission of Academic Press. Copyright © 1993.
- ### Chapter 16
- Figure 16.3: Ulrich, A., Hills, F. J. (1969) *Sugar Beet Nutrient Deficiency Symptoms. A Color Atlas and Chemical Guide*. University of California, Division of Agriculture and Natural Resources, Davis. Copyright © 1969 University of California Board of Regents.
- Figure 16.5: Adapted from Stacey, G., Burris, R. H., Evans, H. J., eds. (1992) *Biological Nitrogen Fixation*. Chapman & Hall, New York. Used with kind permission from Kluwer Academic Publishers.
- Figure 16.6B: B. Newton, Virginia Polytechnic Institute and State University, Blacksburg, Virginia, unpublished; based on data from Kim, J., Rees, D. C. (1994) Ribbon structure of nitrogenase complex including Fe and MoFe proteins. Nitrogenase and biological nitrogen fixation. *Biochemistry* 33: 389–397.
- Figure 16.7: Adapted by permission from Schindelin, H., Kisker, C., Schlessman, J. L., Howard, J. B., Rees, D. C. (1987) Structure of ADP•AIF₄-stabilized nitrogenase complex and its implications for signal transduction. *Nature* 387: 370–376. Copyright © 1987 Macmillan Magazines Ltd.
- Figure 16.8: Peters, J. W., Stowell, M.H.B., Soltis, S. M., Finnegan, M. G., Johnson, M. K., Rees, D. C. (1997) Redox-dependent structural changes in the nitrogenase P-cluster. *Biochemistry* 36: 1181–1187. Copyright © 1997 American Chemical Society.
- Figure 16.10: Reprinted by permission from Schindelin, H., Kisker, C., Schlessman, J. L., Howard, J. B., Rees, D. C. (1987) Structure of ADP•AIF₄-stabilized nitrogenase complex and its implications for signal transduction. *Nature* 387: 370–376. Copyright © 1987 Macmillan Magazines Ltd.
- Figure 16.11: Lowe, D. J. (1992) A role of the 'P' centres of *Klebsiella pneumoniae* MoFe protein in reducing dinitrogen. In *Ninth International Congress on Nitrogen Fixation*, R. Palacios, J. Mora, and W. E. Newton, eds. Kluwer Academic Publishers, Dordrecht, The Netherlands. Used with kind permission from Kluwer Academic Publishers.
- Figure 16.12: L. Feldman, University of California, Berkeley; previously unpublished.
- Figure 16.13: Adapted from Soltis, D. E., Soltis, P. S., Morgan, D. R., Swensen, S. M., Mullin, B. C., Dowd, J. M., Martin, P. G. (1995) Chloroplast gene sequence data suggest a single origin of the predisposition for symbiotic nitrogen fixation in angiosperms. *Proc. Natl. Acad. Sci. USA* 92: 2647–2651. Copyright © 1995 National Academy of Sciences, U.S.A.
- Figure 16.14: S. R. Long, Stanford University, Palo Alto, CA; previously unpublished.
- Figure 16.15: Adapted from van Berkum, P., Eardly, B. D. (1998) Molecular evolutionary systematics of the Rhizobiaceae. In *The Rhizobiaceae*, H. P. Spaink, A. Kondrosi, and P.J.J. Hooykaas, eds. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp. 1–24. Used with kind permission from Kluwer Academic Publishers.
- Figure 16.17A,B: Photographs: D. Ehrhardt and S. R. Long, Stanford University, Palo Alto, CA; previously unpublished.
- Figure 16.17C: Photographs: VandenBosch, K. A., Newcomb, E. H. (1986) Immunogold localization of nodule-specific uricase in developing soybean root nodules. *Planta* 167: 425–436. Copyright © 1987 Springer-Verlag.
- Figure 16.21A: Maps for *R. leuminosarum* bv. *viciae*, *S. meliloti*, and *B. japonicum* adapted from Schlaman, H.R.M., Phillips,

- D. A., Kondorosi, E. (1998) Genetic organization and transcriptional regulation of rhizobial nodulation genes. In *The Rhizobiaceae*, H. P. Spaink, A. Kondorosi, and P.J.J. Hooykaas, eds. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp. 361–386. Used with kind permission from Kluwer Academic Publishers.
- Figure 16.21B: Map for pNGR34 adapted by permission from Freiberg, C., Fellay, R., Bairoch, A., Broughton, W. J., Rosenthal, A., Perret, X. (1997) Molecular basis of symbiosis between *Rhizobium* and legumes. *Nature* 387: 394–401. Copyright © 1997 Macmillan Magazines Ltd.
- Figure 16.25: Adapted from Becker, A., Pühler, A. (1999) Production of exopolysaccharides. In *The Rhizobiaceae*, H. P. Spaink, A. Kondorosi, and P.J.J. Hooykaas, eds. Kluwer Academic Publishers, Dordrecht, The Netherlands, p. 110. Used with kind permission from Kluwer Academic Publishers.
- Figure 16.26: Felle, H. H., Kondorosi, E., Kondorosi, A., Schultze, M. (1998) The role of ion fluxes in Nod factor signalling in *Mendicago sativa*. *Plant J.* 13: 455–463. Reprinted with permission from Blackwell Science Ltd.
- Figure 16.27: Ehrhardt, D. W., Wais, R., Long, S. R. (1996) Calcium spiking in plant root hairs responding to *Rhizobium* nodulation signals. *Cell* 85: 673–681. Copyright © 1996 Cell Press.
- Figure 16.28: Scheres, B., Van De Wiel, C., Zalensky, A., Horvath, B., Spaink, H., Van Eck, H., Zwartkruis, F., Wolters, A.-M., Gloudemans, T., Van Kammen, A., Bisseling, T. (1990) The ENOD12 gene product is involved in the infection process during the pea–*Rhizobium* interaction. *Cell* 60: 281–294. Copyright © 1990 Cell Press.
- Figure 16.29: Heidstra, R., Yang, W. C., Yalcin, Y., Peck, S., Emons, A., Van Kammen, A., Bisseing, T. (1997) Ethylene provides positional information on cortical cell division but is not involved in Nod factor-induced root hair tip growth in *Rhizobium*–legume interaction. *Development* 124: 1781–1787. Used with permission from Company of Biologists Ltd.
- Figure 16.35A,B: Adapted from Touraine, B., Glass, A.D.M. (1997) NO_3^- and ClO_3^- fluxes in the *chl1-5* mutant of *Arabidopsis thaliana*—does the *CHL1-5* gene encode a low-affinity NO_3^- transporter? *Plant Physiol.* 114: 137–144.
- Figure 16.36: N. M. Crawford, University of California–San Diego, La Jolla, CA, previously unpublished.
- Figure 16.37: Adapted from Wang, R., Crawford, N. M. (1996) Genetic identification of a gene involved in constitutive, high affinity, nitrate transport in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* 93: 9297–9301. Copyright © 1996 National Academy of Sciences, U.S.A.
- Figure 16.38B: W. H. Campbell, Michigan Technical Institute, Houghton; previously unpublished.
- Figure 16.39: Campbell, W. H. (1999) Nitrate reductase structure, function and regulation: bridging the gap between biochemistry and physiology. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 50: 277–303. Copyright © 1999 Annual Reviews. www. AnnualReviews.org
- Figure 16.40: Reprinted from Guoguang, L., Campbell, W. H., Schneider, G., Lindquist, Y. (1994) Crystal structure of the FAD-containing fragment of corn nitrate reductase at 2.5 Å resolution: relationship to other flavoprotein reductases. *Structure* 2: 809–821. Copyright © 1994 with permission from Elsevier Science.
- Figure 16.41A: Adapted from Melzer, J. M., Kleinhofs, A., Warner, R. L. (1989) Nitrate reductase regulation: effects of nitrate and light on nitrate reductase mRNA accumulation. *Mol. Gen. Genet.* 217: 341–346. Copyright © 1989 Springer-Verlag.
- Figure 16.41B: Adapted from Pilgrim, M. L., Caspar, T., Quail, P. H., McClung, C. R. (1993) Circadian and light-regulated expression of nitrate reductase in *Arabidopsis*. *Plant Mol. Biol.* 23: 349–364. Used with kind permission from Kluwer Academic Publishers.
- Figure 16.42: Adapted from Huber, S. C., Bachmann, M., Huber, J. L. (1996) Post-translational regulation of nitrate reductase activity: a role for Ca^{2+} and 14-3-3 proteins. *Trends Plant Sci.* 1: 432–438. Copyright © 1996 with permission from Elsevier Science.
- Figure 16.52: Schnug, E., Haneklaus, S. (1994) Symptomatology of S deficiency in *Brassica* species. *Landbauforsch. Voelkenrode Sonderh.* 144: 14–21. Reprinted with permission.
- Figure 16.53: Byers, M., Franklin, J., Smith, S. J. (1987) The nitrogen and sulphur nutrition of wheat and its effect on the composition and baking quality of the grain. *Aspects Appl. Biol.* 15: 337–344. Used by permission from the Association of Applied Biologists.
- Figure 16.57: Adapted from Nissen, P. (1971) Uptake of sulfate by roots and leaf slices of barley mediated by single, multiphasic mechanisms. *Physiol. Plant.* 24: 315–324. Copyright © 1971 Physiologium Plantarum and published by Munksgaard International Publishers Ltd., Copenhagen, Denmark.
- Figure 16.58: Hawkesford, M. J., Smith, F. W. (1997) Molecular biology of higher plant sulphate transporters. In *Sulphur Metabolism in Higher Plants*, W. J. Cram, L. J. DeKok, I. Stulen, C. Brunold, and H. Rennenberg, eds. Backhuys Publishers, Leiden, The Netherlands, pp. 13–26. Used with permission.
- Figure 16.63A,B: Sangman, L. (1999) Molecular analysis of sulfate assimilation in higher plants: effect of cysteine, sulfur, and nitrogen nutrients, heavy metal stress, and genomic DNA cloning. Ph.D. thesis. Rutgers, The State University of New Jersey, New Brunswick, NJ, 155 pp. Used with permission of the author.
- Figure 16.69B: Adapted from Rauser, W. (1995) Phytochelatin and related peptides. *Plant Physiol.* 109: 1141–1149.
- Box 16.1: Chinese characters: Ho, P.-T. (1997) Indigenous origins of Chinese agriculture. In *Origins of Agriculture*, C. A. Reed, ed. Mouton Publishers, The Hague. Flowchart: National Fertilizer Development Center, Tennessee Valley Authority, Chattanooga, TN.
- Table 16.2: Vitousek, P. M., Aber, J. D., Howart, R. W., Likens, G. E., Matson, P. A., Schindler, D. W., Schlesinger, W. H., Tilman, D. G. (1997) Human alteration of the global nitrogen cycle: sources and consequences. *Ecol. Appl.* 7: 737–750. Reprinted with permission from Ecological Society of America.

Chapter 17

Figure 17.5: A. Crozier, University of Glasgow, UK; previously unpublished.

Figure 17.6: Lang, A. (1957) The effect of gibberellin upon flower formation. *Proc. Natl. Acad. Sci. USA* 43: 709–717.

Figure 17.7: A. Crozier, University of Glasgow, UK; previously unpublished.

Figure 17.11A–G: Silverstone, A. L., Chang, C.-W., Krol, E., Sun, T.-P. (1997) Developmental regulation of the gibberellin biosynthetic gene *GAI* in *Arabidopsis thaliana*. *Plant J.* 12: 9–119. Used with permission from Blackwell Science Ltd.

Figure 17.14: Phillips, A. L., Ward, D. A., Ukness, S., Appleford, N.E.J., Lange, T., Huttly, A. K., Gaskin, P., Graebe, J. E., Hedden, P. (1995) Isolation and expression of three gibberellin 20-oxidase cDNA clones from *Arabidopsis*. *Plant Physiol.* 108: 1049–1057.

Figure 17.18: Phillips, A. L., Ward, D. A., Ukness, S., Appleford, N.E.J., Lange, T., Huttly, A. K., Gaskin, P., Graebe, J. E., Hedden, P. (1995) Isolation and expression of three gibberellin 20-oxidase cDNA clones from *Arabidopsis*. *Plant Physiol.* 108: 1049–1057.

Figure 17.19: Adapted from Gilmour, S. J., Zeevaart, J.A.D., Schwenen, L., Graebe, J. E. (1986) Gibberellin metabolism in cell-free extracts from spinach (*Spinacia oleracea*) leaves in relation to photoperiod. *Plant Physiol.* 82: 190–195.

Figure 17.20: Electrophoretograms: Toyomasu, T., Kawaide, H., Mitsushashi, W., Inoue, Y., Kamiya, Y. (1998) Phytochrome regulates gibberellin biosynthesis during germination of photoblastic lettuce seeds. *Plant Physiol.* 118: 1517–1523.

Figure 17.22: S. McCormick, University of California, Berkeley; previously unpublished.

- Figure 17.23: J. Weyers, University of Dundee, UK; previously unpublished.
- Figure 17.31: T. Kakimoto, Osaka University, Japan; previously unpublished.
- Figure 17.33: I. Scott, University of Wales, Aberystwyth, UK; previously unpublished.
- Figure 17.34: G. Sandberg, Swedish University of Agricultural Sciences, Umeå; previously unpublished.
- Figure 17.35A,B: Faiss, M., Zalubilova, J., Strand, M., Schmülling, T. (1997) Conditional transgenic expression of the *ipt* gene indicates a function for cytokinins in paracrine signaling in whole tobacco plants. *Plant J.* 12: 401–415. Used with permission from Blackwell Science Ltd.
- Figure 17.36: Martineau, B., Houck, C. M., Sheehy, R. E., Hiatt, W. R. (1994) Fruit-specific expression of the *A. tumefaciens* isopentenyl transferase gene in tomato: effects on fruit ripening and defense-related gene expression in leaves. *Plant J.* 5: 11–19. Used with permission from Blackwell Science Ltd.
- Figure 17.43: J. Cohen, U.S. Department of Agriculture, Beltsville, MD, and A. Wright, University of Missouri, Columbia; previously unpublished.
- Figure 17.49: Adapted from Normanly, J., Grisafi, P., Fink, G. R., Bartel, B. (1997) *Arabidopsis* mutants resistant to the auxin effects of indole-3-acetonitrile are defective in the nitrilase encoded by the *NIT1* gene. *Plant Cell* 9: 1781–1790.
- Figure 17.51: Adapted from Nilsson, O., Crozier, A., Schmülling, T., Sandberg, G., Olsson, O. (1993) Indole-3-acetic acid homeostasis in transgenic tobacco plants expressing *Agrobacterium rhizogenes rolB* gene. *Plant J.* 3: 681–689. Used with permission from Blackwell Science Ltd.
- Figure 17.52: Romano, C. P., Cooper, M. L., Klee, H. J. (1993) Uncoupling auxin and ethylene effects in transgenic tobacco and *Arabidopsis* plants. *Plant Cell* 5: 181–189.
- Figure 17.53A–C: H. Mori, Nagoya University, Japan; previously unpublished.
- Figure 17.56: D. Grierson, University of Nottingham, UK; previously unpublished.
- Figure 17.58: T. Yokota, Teikyo University Utsunomiya, Japan; previously unpublished.
- Figure 17.60: T. Yokota, Teikyo University, Utsunomiya, Japan; previously unpublished.
- Figure 17.71: Adapted from Kumar, A., Altabella, T., Taylor, M. A., Tiburcio, A. F. (1997) Recent advances in polyamine research. *Trends Plant Sci.* 2: 124–130. Copyright © 1997 with permission from Elsevier Science.
- Figure 17.74 A,B: Kubigsteltig, I., Laudert, D., Weiler, E. W. (1999) Structure and regulation of the *Arabidopsis thaliana* allene oxidase synthase gene. *Planta* 208: 463–471. Copyright © 1999 Springer-Verlag.
- Figure 17.78: Reprinted with permission from Raskin, I., Ehmann, A., Melander, W. R., Meeuse, B.J.D. (1987) Salicylic acid—a natural inducer of heat production in Arum lilies. *Science* 237: 1545–1556. Copyright © 1987 American Association for the Advancement of Science.
- Figure 17.79: J. Draper, University of Wales, Aberystwyth, UK; previously unpublished.
- Figure 17.81: J. Draper, University of Wales, Aberystwyth, UK; previously unpublished.
- Table 17.2: Kumar, A., Taylor, M. A., Arif, S.A.M., Davies, H. V. (1996) Potato plants expressing antisense and sense *S*-adenosylmethionine decarboxylase (SAMDC) transgenes show altered levels of polyamines and ethylene: antisense plants display abnormal phenotypes. *Plant J.* 9: 147–158. Used with permission from Blackwell Science Ltd.
- ### Chapter 18
- Figure 18.2: Staehelin, L. A., Giddings, T. H. Jr., Kiss, J. Z., Sack, F. D. (1990) Macromolecular differentiation of Golgi stacks in root tips of *Arabidopsis* and *Nicotiana* seedlings as visualized in high-pressure frozen and freeze-substituted samples. *Protoplasma* 157: 75–91. Copyright © 1990 Springer-Verlag.
- Figure 18.4: X.-W. Deng, Yale University, New Haven, CT; previously unpublished.
- Figure 18.7: Adapted from Wallace, D.H., Enriquez, G.A. (1980) Daylength and temperature effects on days to flowering of early and late maturing beans (*Phaseolus vulgaris* L.). *J. Am. Hort. Sci.* 105: 583–591.
- Figure 18.9: W. Gruissem, University of California, Berkeley; previously unpublished.
- Figure 18.14: Adapted from Plakidou-Dymock, S., Dymock, D., Hooley, R. (1998) A higher plant seven-transmembrane receptor that influences sensitivity to cytokinins. *Curr. Biol.* 8: 315–324. Copyright © 1998 with permission from Elsevier Science.
- Figure 18.22: Wilkinson, J. Q., Lanahan, M. B., Clark, D. G., Bleecker, A. B., Chang, C., Meyerowitz, E. M., Klee, H. J. (1997) A dominant mutant receptor from *Arabidopsis* confers ethylene insensitivity in heterologous plants. *Nat. Biotechnol.* 15: 444–447.
- Figure 18.25: A. J. Trewavas, Institute of Cell and Molecular Biology, University of Edinburgh, UK; previously unpublished.
- Figure 18.26A,B: Data from Allan, A. C., Fricker, M. D., Ward, J. L., Beale, M. H., Trewavas, A. J. (1994) Two transduction pathways mediate rapid effects of abscisic acid in *Commelina* guard cells. *Plant Cell* 6: 1319–1328.
- Figure 18.30: Adapted from Wittinghofer, A., Pai, E. F. (1991) The structure of Ras protein: a model for a universal molecular switch. *Trends Biochem. Sci.* 16: 382–387.
- Copyright © 1991 with permission from Elsevier Science.
- Figure 18.36: Adapted from Munnik, T., Irvine, R. F., and Musgrave, A. (1998) Phospholipid signalling in plants. *Biochim. Biophys. Acta* 1389: 222–272.
- Figure 18.37: Adapted from Munnik, T., Irvine, R. F., and Musgrave, A. (1998) Phospholipid signalling in plants. *Biochim. Biophys. Acta* 1389: 222–272.
- Figure 18.46: A. J. Trewavas, Institute of Cell and Molecular Biology, University of Edinburgh, Scotland; previously unpublished.
- Figure 18.47: Reprinted by permission from Shacklock, P. S., Read, N. D., Trewavas, A. J. (1992) Cytosolic free calcium mediates red light-induced photomorphogenesis. *Nature* 358: 753–755. Copyright © 1992 Macmillan Magazines Ltd.
- Figure 18.48: Mahlo, R., Read, N. D., Pais, M. S., Trewavas, A. J. (1994) Role of cytosolic free calcium in the reorientation of pollen tube growth. *Plant J.* 5: 331–341. Used with permission from Blackwell Science Ltd.
- Figure 18.51: A. J. Trewavas, Institute of Cell and Molecular Biology, University of Edinburgh, Scotland; previously unpublished.
- Figure 18.52: Gilroy, S., Fricker, M. D., Read, N. D., Trewavas, A. J. (1991) Role of calcium in signal transduction of *Commelina* guard cells. *Plant Cell* 3: 333–341.
- Figure 18.53B: Pierson, E. S., Miller, D. D., Callahan, D. A., Van Aken, J., Hackett, G., Hepler, P. K. (1996) Tip-localized calcium entry fluctuates during pollen tube growth. *Dev. Biol.* 174: 160–173. Reprinted by permission of Academic Press. Copyright © 1996.
- Figure 18.54A: Franklin-Tong, V. E., Drobak, B. K., Allan, A. C., Watkins, P.A.C., Trewavas, A. J. (1996) Growth of pollen tubes of *Papaver rhoeas* is regulated by a slow-moving calcium wave propagated by inositol 1,4,5-trisphosphate. *Plant Cell* 8: 1305–1321.
- Figure 18.55: Reproduced from Knight, M. R., Read, N. D., Campbell, A. K., Trewavas, A. J. (1993) Imaging calcium dynamics in living plants using semi-synthetic recombinant aequorins. *J. Cell Biol.* 121: 83–90. By copyright permission from the Rockefeller University Press.
- Figure 18.57: Adapted with permission from Ikura, M., Clore, G. M., Gronenborn, A. M., Zhu, G., Klee, C. B., Bax, A. (1992) Solution structure of a calmodulin-target peptide complex by multidimensional NMR. *Science* 256: 632–638. Published by American Association for the Advancement of Science.
- Figure 18.65A: K. Stepnitz, Michigan State University, East Lansing; published as the cover of *Science*, 26 August 1988,

accompanying the paper by Bleecker, A. B., Estelle, M. A., Somerville, C., Kende, H. (1988) Insensitivity to ethylene conferred by a dominant mutation in *Arabidopsis thaliana*. *Science* 241: 1086–1089. Used with permission of the photographer.

Figure 18.65B: J. Keiber, University of Chicago; previously unpublished.

Fig. 18.73: Adapted with permission from DeKoninck, P., Schulman, H. (1998) Sensitivity of CaM kinase II to the frequency of Ca²⁺ oscillations. *Science* 279: 227–230. Copyright © 1998 American Association for the Advancement of Science.

Box 18.4: Adapted from Furuya, M., Schaffer, E. (1996) Photo perception and signaling of induction reactions by different phytochromes. *Trends Plant Sci.* 1: 301–307. Copyright © 1996 by permission from Elsevier Science.

Box 18.5A,B: Mahlo, R., Trewavas, A. J. (1996) Localized apical increases of cytosolic free calcium control pollen tube orientation. *Plant Cell* 8: 1935–1949.

Box 18.6B,C: Data from Trewavas, A. J., Malho, R. (1998) Ca²⁺ signalling in plant cells: the big network! *Curr. Opin. Plant Biol.* 1: 428–434.

Box 18.6D: H. Page, N. D. Read, and A. J. Trewavas, Institute of Cell and Molecular Biology, University of Edinburgh, Scotland; previously unpublished.

Table 18.1: Data from Hewitt, E. J. (1963) Inorganic nutrition. In *Plant Physiology; a Treatise*, Vol. 3. F. C. Steward, ed. Academic Press, New York, pp. 137–361.

Chapter 19

Figure 19.1: Adapted from Goldberg, R. B. (1988) Plants' novel developmental processes. *Science* 240: 1460–1467.

Figure 19.3: Adapted from Lang, A. (1965) Physiology of flower initiation. In *Handbuch der Pflanzenphysiologie (Encyclopedia of Plant Physiology)*, W. Ruhland, ed. Springer-Verlag, Berlin, pp. 1379–1536. Copyright © 1965 Springer-Verlag.

Figure 19.4: F. Hempel, University of California, Berkeley; previously unpublished.

Figure 19.5: G. Coupland, John Innes Centre, Norwich, UK; previously unpublished.

Figure 19.7A,B: R. Amasino, University of Wisconsin, Madison; previously unpublished.

Figure 19.8: D. Weigel, Salk Institute, San Diego, CA; previously unpublished.

Figure 19.9A: Photograph: Shannon, S., Meeks-Wagner, D. R. (1991) A mutation in the *Arabidopsis TFL1* gene affects inflorescence meristem development. *Plant Cell* 3: 877–892.

Figure 19.9B: Adapted with permission from Bradley, D., Ratcliffe, O., Vincent, C., Carpenter, R., Coen, E. (1997) Inflorescence

commitment and architecture in *Arabidopsis*. *Science* 275: 80–83. Copyright © 1997 American Association for the Advancement of Science.

Figure 19.12A,B: Smyth, D. R., Bowman, J. L., Meyerowitz, E. M. (1990) Early flower development in *Arabidopsis*. *Plant Cell* 2: 755–767.

Figure 19.13: Smyth, D. R., Bowman, J. L., Meyerowitz, E. M. (1990) Early flower development in *Arabidopsis*. *Plant Cell* 2: 755–767.

Figure 19.14: Smyth, D. R., Bowman, J. L., Meyerowitz, E. M. (1990) Early flower development in *Arabidopsis*. *Plant Cell* 2: 755–767.

Figure 19.15: Meyerowitz, E. M., Bowman, J. L., Brockman, L. L., Drews, G. N., Jack, T., Sieburth, L. E., Weigel, D. (1991) A genetic and molecular model for flower development in *Arabidopsis thaliana*. *Development (Suppl. 1)*: 157–167. Used with permission from Company of Biologists Ltd.

Figure 19.18A: Adapted from Okamoto, J. K., den Boer, B.G.W., Jofuku, K. D. (1993) Regulation of *Arabidopsis* flower development. *Plant Cell* 5: 1183–1193.

Figure 19.19: Münster, T., Pahnke, J., Di Rosa, A., Kim, J. T., Martin, W., Saedler, H., Theissen, G. (1997) Floral homeotic genes were recruited from homologous MADS-box genes preexisting in the common ancestor of ferns and seed plants. *Proc. Natl. Acad. Sci. USA* 94: 2415–2420. Copyright © 1997 National Academy of Sciences, U.S.A.

Figure 19.20: Chen, L., Cheng, J.-C., Castle, L., Sung, Z. R. (1997) *EMF* genes regulate *Arabidopsis* inflorescence development. *Plant Cell* 9: 2011–2024.

Figure 19.22: Wagner, V. T., Cresti, M., Salvatice, P., Tiezzi, A. (1990) Changes in volume, surface area, and frequency of nuclear pores on the vegetative nucleus of tobacco pollen in fresh, hydrated, and activated conditions. *Planta* 181: 304–309. Copyright © 1990 Springer-Verlag.

Figure 19.23: Plantain, buttercup, ox-eye daisy, nettle: M. P. Kage, Okapia, Frankfurt, Germany; published in Attenborough, D. (1995) *The Private Lives of Plants. A Natural History of Plant Behaviour*. Princeton University Press, Princeton, NJ. Used with permission from Okapia. Winter's bark, morning glory, rose cactus, aheaahea: Gifford, E. M., and Foster, A. S. (1987) *Morphology and Evolution of Vascular Plants*, 3rd ed. W. H. Freeman and Co., New York. Used with permission from E. M. Gifford. Poppy: Stern, K. R. (1991) *Introductory Plant Biology*, 5th ed. W. C. Brown Publishers, Dubuque, IA. Used with permission from K. R. Stern. Ragweed: Howlett, B. J., Knox, R. B., Heslop-Harrison, J. (1973) Pollen-wall proteins: release of the allergen antigen E from intine and exine sites in pollen grains of ragweed and *Cosmos*. *J. Cell Sci.* 13: 603–619. Used with permission from Company of Biologists Ltd.

Sycamore and hollyhock: Andrew & Polly Syred, Microscopix, Powys, UK. Used with permission from Microscopix.

Figure 19.24: Photograph: Drews, G. N., Lee, D., Christensen, C. (1998) Genetic analysis of female gametophyte development. *Plant Cell* 10: 5–17.

Figure 19.25: Adapted from Nadeau, J. A., Zhang, X. S., Li, J., O'Neill, S. D. (1996) Ovule development: identification of state-specific and tissue-specific cDNAs. *Plant Cell* 8: 213–239.

Figure 19.26: Drews, G. N., Lee, D., Christensen, C. (1998) Genetic analysis of female gametophyte development. *Plant Cell* 10: 5–17.

Figure 19.27: U. Grossniklaus, Basel, Switzerland; published as the cover of the Cold Spring Harbor Laboratory 1998 Annual Report, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.

Figure 19.28: Adapted from Wise, R. P., Brownson, C., Schnable, P. S., Horner, H. T. (1998) T-cytoplasmic male sterility of maize. *Adv. Agron.* 65: 79–130. Reprinted by permission of Academic Press. Copyright © 1998.

Figure 19.29A: Adapted from Mascarenhas, J. P. (1993) Molecular mechanisms of pollen tube growth and differentiation. *Plant Cell* 5: 1303–1314.

Figure 19.29B: Photograph: Newbigin, E., Anderson, M. A., Clarke, A. E. (1993) Gametophytic self-incompatibility systems. *Plant Cell* 5: 1315–1324.

Figure 19.30: Jauh, G. Y., Eckhard, K. J., Nothnagal, E. A., Lord, E. M. (1997) Adhesion of lily pollen tubes on an artificial matrix. *Sex. Plant Reprod.* 10: 173–180. Copyright © 1997 Springer-Verlag.

Figure 19.31A–D: Higashiyama, T., Kuroiwa, H., Kawano, S., Kuroiwa, T. (1998) Guidance in vitro of the pollen tube to the naked embryo sac of *Torenia fournieri*. *Plant Cell* 10: 2019–2031.

Figure 19.32: Adapted from McCormick, S. (1998) Self-incompatibility and other pollen-pistil interactions. *Curr. Opin. Plant Biol.* 1: 18–25.

Figure 19.33: Photograph: Nasrallah, J. B., Nasrallah, M. E. (1993) Pollen-stigma signaling in the sporophytic self-incompatibility response. *Plant Cell* 5: 1325–1335.

Figure 19.34A–D: Photograph: Dickinson, H. G., Elleman, C. J. (1994) Pollen hydrodynamics and self-incompatibility in *Brassica oleracea*. In *Pollen–Pistil Interactions and Pollen Tube Growth (Current Topics in Plant Physiology)*, Vol. 12. A. G. Stephenson and T.-H. Kao, eds. American Society of Plant Physiologists, Rockville, MD, pp. 45–61.

Figure 19.36: Photographs and drawings are from the same respective sources: Preglobular through heart—Yadegari, R., de Paiva, G. R., Laux, T., Koltunow, A. M., Apuya, N., Zimmerman, J. L., Fischer, R. L., Harada, J. J., Goldberg, R. B. (1994) Cell

differentiation and morphogenesis are uncoupled in *Arabidopsis* raspberry embryos. *Plant Cell* 6: 1713–1729. Torpedo and mature embryo—West, M. A., Harada, J. J. (1993) Embryogenesis in higher plants: an overview. *Plant Cell* 5: 1361–1369.

Figure 19.37: Photographs: leafy cotyledon and fusca—West, M. A., Yee, K. M., Danao, J., Zimmerman, J. L., Fischer, R. L., Goldberg, R. B., Harada, J. J. (1994) *LEAFY COTYLEDON1* is an essential regulator of late embryogenesis and cotyledon identity in *Arabidopsis*. *Plant Cell* 6: 1731–1745. raspberry—Yadegari, R., de Paiva, G. R., Laux, T., Koltunow, A. M., Apuya, N., Zimmerman, J. L., Fischer, R. L., Harada, J. J., Goldberg, R. B. (1994) Cell differentiation and morphogenesis are uncoupled in *Arabidopsis* raspberry embryos. *Plant Cell* 6: 1713–1729.

Figure 19.38: Lopes, M. A., Larkins, B. A. (1993) Endosperm origin, development, and function. *Plant Cell* 5: 1383–1399.

Figure 19.45: Adapted from Krochko, J. E., Bewley, J. D. (1989) Use of electrophoretic techniques in determining the composition of seed storage proteins in alfalfa. *Electrophoresis* 9: 751–763. Copyright © 1989 Springer-Verlag.

Figure 19.47: Data from Gatehouse, J. A., Evans, I. M., Croy, R.R.D., Boulter, D. (1986) Differential expression of genes during legume seed development. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 314: 367–384; and Boulter, D., Evans, I. M., Ellis, J. R., Shirsat, A., Gatehouse, J. A., Croy, R.R.D. (1987) Differential gene expression in the development of *Pisum-sativum*. *Plant Physiol. Biochem. (Paris)* 25: 283–290. Copyright Elsevier Science.

Figure 19.48: Adapted from Kawagoe, Y., Murai, N. (1992) Four distinct nuclear proteins recognize in vitro the proximal promoter of the bean seed storage protein β -phaseolin conferring spatial and temporal control. *Plant J.* 2: 927–936; and data from Burrow, M. D., Sen, P., Chlan, C. A., Murai, N. (1992) Developmental control of the β -phaseolin gene requires positive, negative, and temporal seed-specific transcriptional regulatory elements and a negative element for stem and root expression. *Plant J.* 2: 537–548.

Figure 19.52: Adapted from Bewley, J. D. (1997) Seed germination and dormancy. *Plant Cell* 9: 1055–1066.

Box 19.2A,B: Gustafson-Brown, C., Savidge, B., Yanofsky, M. F. (1994) Regulation of the *Arabidopsis* floral homeotic gene *APETA-LA1*. *Cell* 76: 131–143. Copyright © 1994 Cell Press.

Box 19.2C: D. Weigel, Salk Institute, San Diego, CA; previously unpublished.

Box 19.2D: Meyerowitz, E. M., Bowman, J. L., Brockman, L. L., Drews, G. N., Jack, T., Sieburth, L. E., Weigel, D. (1991) A genetic and molecular model for flower develop-

ment in *Arabidopsis thaliana*. *Development* (Suppl. 1): 157–167. Used with permission from Company of Biologists Ltd.

Box 19.2E,H: Zachgo, S., Silva, E. de A., Motte, P., Tröbner, W., Saedler, H., Schwarz-Sommer, Z. (1995) Functional analysis of the *Antirrhinum* floral homeotic *DEFICIENS* gene in vivo and in vitro by using a temperature-sensitive mutant. *Development* 121: 2861–2875. Used with permission from Company of Biologists Ltd.

Box 19.2F: Huijser, P., Klein, J., Lönning, W.-E., Meijer, H., Saedler, H., Sommer, H. (1992) Bracteomania, an inflorescence anomaly, is caused by the loss of function of the *MADS*-box gene *squamosa* in *Antirrhinum majus*. *EMBO J.* 11: 1239–1249. Used with permission from Oxford University Press.

Box 19.2G: Carpenter, R., Copsey, L., Vincent, C., Doyle, S., Magrath, R., Coen, E. (1995) Control of flower development and phyllotaxy by meristem identity genes in *Antirrhinum*. *Plant Cell* 7: 2001–2011.

Box 19.3: Adapted from McCormick, S. (1991) Molecular analysis of male gametogenesis in plants. *Trends Genet.* 7: 298–303. Copyright © 1991 with permission from Elsevier Science.

Box 19.6A,B: Lolle, S. J., Cheung, A. Y. (1993) Promiscuous germination and growth of wildtype pollen from *Arabidopsis* and related species on the shoot of the *Arabidopsis* mutant, *fiddlehead*. *Dev. Biol.* 155: 250–258. Reprinted by permission of Academic Press. Copyright © 1993.

Box 19.7: Kranz, E., Bautor, J., Lörz, H. (1994) In vitro fertilization of single, isolated gametes of maize mediated by electrofusion. *Sex. Plant Reprod.* 4: 12–16.

Table 19.1: Smyth, D. R., Bowman, J. L., Meyerowitz, E. M. (1990) Early flower development in *Arabidopsis*. *Plant Cell* 2: 755–767.

Table 19.2: Data from Bewley, J. D., and Black, M. (1994) *Seeds. Physiology of Development and Germination*. Plenum Press, New York.

Chapter 20

Figure 20.2A: Reproduced from Takeshige, K., Baba, M., Tsuboi, S., Noda, T., Ohsumi, Y. (1992) Autophagy in yeast demonstrated with proteinase-deficient plants and conditions for its induction. *J. Cell Biol.* 119: 301–311. By copyright permission of the Rockefeller University Press.

Figure 20.2B: Adapted from Matile, P., Winklenbach, F. (1971) Function of lysosomes and lysosomal enzymes in the senescing corolla of the morning glory (*Ipomoea purpurea [sic]*). *J. Exp. Bot.* 22: 759–771. Used with permission from Oxford University Press.

Figure 20.3: P. Bethke, R. Jones, University of California, Berkeley; previously unpublished.

Figure 20.5A: Neuffer, M. G., Coe, E. H., Wessler, S. R. (1997) *Mutants of Maize*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. Copyright © 1997 Cold Spring Harbor Laboratory Press.

Figure 20.5B: L. Jesaitis, University of California, Berkeley; previously unpublished.

Figure 20.6: R. A. Dietrich, Novartis Crop Protection, Inc., Research Triangle Park, NC; previously unpublished.

Figure 20.8A,B: He, C.-J., Morgan, P. W., Drew, M. C. (1996) Transduction of an ethylene signal is required for cell death and lysis in the root cortex of maize during aerenchyma formation induced by hypoxia. *Plant Physiol.* 112: 463–472.

Figure 20.9: Bleeker, A. B., Patterson, S. E. (1997) Last exit: senescence, abscission, and meristem arrest in *Arabidopsis*. *Plant Cell* 9: 1169–1179.

Figure 20.10: Kende, H., Baumgartner, B. (1974) Regulation of aging in flower of *Ipomoea tricolor* by ethylene. *Planta* 116: 279–289. Copyright © 1974 Springer-Verlag.

Figure 20.11: H. Thomas, IGER, Aberystwyth, UK, previously unpublished.

Figure 20.12: Adapted from Noodèn, L. D., Guimet, J. J., John, I. (1997) Senescence mechanisms. *Physiol. Plant.* 101: 746–753. Copyright © 1997 Physiologium Plantarum and published by Munksgaard International Publishers Ltd., Copenhagen, Denmark.

Figure 20.13A–C: Thomas, H. (1977) Ultrastructure, polypeptide composition and photochemical activity of chloroplasts during foliar senescence of a non-yellowing mutant genotype of *Festuca pratensis*. *Planta* 137: 53–60. Copyright © 1977 Springer-Verlag.

Figure 20.14: Adapted from Buchanan-Wollaston, V. (1997) The molecular biology of leaf senescence. *J. Exp. Bot.* 48: 181–199. Reprinted with permission from Oxford University Press.

Figure 20.16: Adapted with permission from Matile, P., Hörtensteiner, S., Thomas, H. (1999) Chlorophyll degradation. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 50: 67–95. Copyright © 1999 Annual Reviews. www.AnnualReviews.org

Figure 20.19A–C: PaO data: Vicentini, F., Hörtensteiner, S., Schellenberg, M., Thomas, H., Matile, P. (1995) Chlorophyll breakdown in senescent leaves: identification of the biochemical lesion in a *stay-green* genotype of *Festuca pratensis* Huds. *New Phytol.* 129: 247–252. Reprinted by permission of Academic Press. Copyright © 1995. Protein data: Adapted from Hilditch, P., Thomas, H., Thomas, B. J., Rogers, L. J. (1989) Leaf senescence in a non-yellowing mutant of *Festuca pratensis*: proteins of Photosystem II. *Planta* 177: 265–272. Copyright © 1997 Springer-Verlag.

Figure 20.20A,B: Griffiths, C. M., Hosken, S. E., Oliver, D., Chojcecki, J., Thomas, H.

(1997) Sequencing, expression pattern and RFLP mapping of a senescence-enhanced cDNA from *Zea mays* with high homology to oryzain γ and aleurain. *Plant Mol. Biol.* 34: 815–821. Used with kind permission of Kluwer Academic Publishers.

Figure 20.20C: I. Donnison, IGER, Aberystwyth, UK, previously unpublished.

Figure 20.23: Adapted from Buchanan-Wollaston, V. (1997) The molecular biology of leaf senescence. *J. Exp. Bot.* 48: 181–199. Reprinted with permission from Oxford University Press.

Figure 20.25: Adapted from Ashihara, H., Takasawa, Y., Suzuki, T. (1997) Metabolic fate of guanosine in higher plants. *Physiol. Plant.* 100: 909–916. Copyright © 1997 Physiologium Plantarum and published by Munksgaard International Publishers Ltd., Copenhagen, Denmark.

Figure 20.26: A. Bleeker, University of Wisconsin, Madison; previously unpublished.

Figure 20.27: Whitehead, C. S., Halevy, A. H., Reid, M. S. (1984) Roles of ethylene and 1-aminocyclopropane-1-carboxylic acid in pollination and wound-induced senescence of *Petunia hybrida* flowers. *Physiol. Plant.* 61: 643–648. Copyright © 1984 Physiologium Plantarum and published by Munksgaard International Publishers Ltd., Copenhagen, Denmark.

Figure 20.28A,B: Lincoln, J. E., Cordes, S., Read, E., Fischer, R. L. (1987) Regulation of gene expression by ethylene during *Lycopersicon esculentum* (tomato) fruit development. *Proc. Natl. Acad. Sci. USA* 84: 2793–2797. Used with permission.

Figure 20.29A–E: Lanahan, M. B., Yen, H. C., Giovannoni, J. J., Klee, H. J. (1994) The *never ripe* mutation blocks ethylene perception in tomato. *Plant Cell* 6: 521–530.

Figure 20.30: Adapted with permission from Oeller, P. W., Min-Wong, L., Taylor, L. P., Pike, D. A., Theologis, A. (1991) Reversible inhibition of tomato fruit senescence by antisense RNA. *Science* 254: 437–439. Copyright © 1991 American Association for the Advancement of Science.

Figure 20.31: Theologis, A., Oeller, P. W., Wong, L. M., Rottmann, W. H., Gantz, D. M. (1993) Use of a tomato mutant constructed with reverse genetics to study fruit ripening, a complex developmental process. *Dev. Genet.* 14: 282–295. Reprinted by permission of Wiley-Liss, Inc., a subsidiary of John Wiley & Sons, Inc.

Figure 20.32: Johal, G. S., Hulbert, S., and Briggs, S. P. (1995) Disease lesion mimics in maize: a model for cell death in plants. *BioEssays* 17: 685–692. Reprinted by permission of Wiley-Liss, Inc., a subsidiary of John Wiley & Sons, Inc.

Figure 20.33: Photograph: Gan, S., Amasino, R. M. (1997) Making sense of senescence. *Plant Physiol.* 113: 313–319.

Figure 20.34: Adapted from Thomas, H., Ougham, H., Hörtensteiner, S. (1998) Ring in the new, ring out the old. *IGER Innovations* 2: 5–9.

Figure 20.35: Raven, P. H. (1999) Cells and tissues of the plant body. In *Biology of Plants*, P. H. Raven, R. F. Evert, and S. E. Eichorn, eds. Worth Publishers, New York, pp. 570–588. Copyright © 1971, 1976, 1981, 1992, and 1999 by Worth Publishers. Used with permission.

Figure 20.36A–F: Adapted from Fukuda, H. (1997) Tracheary element differentiation. *Plant Cell* 9: 1147–1156.

Figure 20.36G,H: A. Groover, University of North Carolina, Chapel Hill; previously unpublished.

Figure 20.37A,B: Adapted from Fukuda, H. (1997) Tracheary element differentiation. *Plant Cell* 9: 1147–1156.

Figure 20.38: Watanabe, Y., Fukada, H. (1995) *Plant Cell Physiol.* (Suppl.): 87. Reprinted with permission from the Japanese Society of Plant Physiologists.

Figure 20.39: Yamamoto, R., Demura, T., Fukuda, H. (1997) Brassinosteroids induce entry into the final stage of tracheary element differentiation in cultured *Zinnia* cells. *Plant Cell Physiol.* 38: 980–983. Reprinted with permission from the Japanese Society of Plant Physiologists.

Figure 20.40A–F: Young, T. E., Gallie, D. R., DeMason, D. A. (1997) Ethylene-mediated programmed cell death during maize endosperm development of wild-type and *shrunken2* genotypes. *Plant Physiol.* 115: 737–751.

Figure 20.41A,B: Young, T. E., Gallie, D. R., DeMason, D. A. (1997) Ethylene-mediated programmed cell death during maize endosperm development of wild-type and *shrunken2* genotypes. *Plant Physiol.* 115: 737–751.

Figure 20.42: Bethke, P. C., Lonsdale, J. E., Fath, A., Jones, R. L. (1999) Hormonally regulated programmed cell death in barley aleurone cells. *Plant Cell* 11: 1033–1045.

Figure 20.43A,B: Bethke, P. C., Lonsdale, J. E., Fath, A., Jones, R. L. (1999) Hormonally regulated programmed cell death in barley aleurone cells. *Plant Cell* 11: 1033–1045.

Figure 20.44A,B: He, C.-J., Drew, M. C., Morgan, P. W. (1994) Induction of enzymes associated with lysigenous aerenchyma formation in roots of *Zea mays* during hypoxia or nitrogen starvation. *Plant Physiol.* 105: 861–865.

Figure 20.45: Subbaiah, C. C., Bush, D. S., Sachs, M. M. (1994) Elevation of cytosolic calcium precedes anoxic gene expression in maize suspension-cultured cells. *Plant Cell* 6: 1747–1762.

Figure 20.46A–C: He, C.-J., Morgan, P. W., Drew, M. C. (1996) Transduction of an ethylene signal is required for cell death and ly-

sis in the root cortex of maize during aerenchyma formation induced by hypoxia. *Plant Physiol.* 112: 463–472.

Figure 20.47: Ward, E. R., Uknes, S. J., Williams, S. C., Dincher, S. S., Wiederhold, D. L., Alexander, D. C., Ahl-Goy, P., Mettraux, J.-P., Ryals, J. A. (1991) Coordinate gene activity in response to agents that induce systemic acquired resistance. *Plant Cell* 3: 1085–1094.

Figure 20.48: Van den Ackerveken, G.F.J.M., Van Kan, J.A.L., De Wit, P.J.G.M. (1992) Molecular analysis of the avirulence gene *avr9* of the fungal tomato pathogen *Cladosporium fulvum* fully supports the gene-for-gene hypothesis. *Plant J.* 2: 359–366. Used with permission from Blackwell Science Ltd.

Figure 20.49A–G: Dangl, J. L., Dietrich, R. A., Richberg, M. H. (1996) Death don't have no mercy: cell death programs in plant-microbe interactions. *Plant Cell* 8: 1793–1807.

Figure 20.50: Mittler, R., Shulaev, V., Lam, E. (1995) Coordinated activation of programmed cell death and defense mechanisms in transgenic tobacco plants expressing a bacterial proton pump. *Plant Cell* 7: 29–42.

Figure 20.51A,B: Freialdenhoven, A., Peterhänsel, C., Kurth, J., Kreuzaler, F., Schulze-Lefert, P. (1996) Identification of genes required for the function of non-race-specific *mlo* resistance to powdery mildew in barley. *Plant Cell* 8: 5–14.

Figure 20.52: Wolter, M., Hollricher, K., Salamini, F., Schulze-Lefert, P. (1993) The *mlo* resistance alleles to powdery mildew infection in barley trigger a developmentally controlled defence mimic phenotype. *Mol. Gen. Genet.* 239: 122–128. Copyright © 1993 Springer-Verlag.

Figure 20.53: Kosslak, R. M., Chamberlin, M. A., Palmer, R. G., Bowen, B. A. (1997) Programmed cell death in the root cortex of soybean *root necrosis* mutants. *Plant J.* 11: 729–745. Used with permission from Blackwell Science Ltd.

Figure 20.54: Adapted from Hammond-Kosack, K. E., Jones, J.D.G. (1996) Resistance gene-dependent plant defense responses. *Plant Cell* 8: 1773–1791.

Figure 20.55: Adapted from Mehdy, M. (1994) Active oxygen species in plant defense against pathogens. *Plant Physiol.* 105: 457–472.

Figure 20.56: Ryerson, D. E., Heath, M. C. (1996) Cleavage of nuclear DNA into oligonucleosomal fragments during cell death induced by fungal infection or by abiotic treatments. *Plant Cell* 8: 393–402.

Figure 20.58: Wang, H., Li, J., Bostock, R. M., Gilchrist, D. G. (1996) Apoptosis: a functional paradigm for programmed plant cell death induced by a host-selective phytotoxin and invoked during development. *Plant Cell* 8: 375–391.

Box 20.1: Gel photograph: Peitsch, M. C., Polzar, B., Stephan, H., Crompton, T., MacDonald, H. R., Mannherz, H. G., Tschopp, J. (1993) Characterization of the endogenous deoxyribonuclease involved in nuclear DNA degradation during apoptosis (programmed cell death). *EMBO J.* 12: 371–377. Reprinted with permission from Oxford University Press.

Table 20.1: Data from Buchanan-Wollaston, V. (1997) The molecular biology of leaf senescence. *J. Exp. Bot.* 48: 181–199. Copyright Oxford University Press.

Part 5 opening photograph:

Galen Rowell/Mountain Light Photography, Emeryville, CA.

Chapter 21

Figure 21.1: K. Hammond-Kosack, Monsanto Co., St. Louis, MO; previously unpublished.

Figure 21.2: E. P. Caswell-Chen, University of California, Davis; published in Moffat, A. S. (1997) First nematode-resistance gene found. *Science* 275: 757.

Figure 21.3: Adapted from Jackson, A. O., Taylor, C. B. (1996) Plant–microbe interactions: life and death at the interface. *Plant Cell* 8: 1651–1668.

Figure 21.4: Reprinted with the permission of Simon & Schuster, from *The Simon & Schuster's Beginner's Guide to Understanding Wine* by M. Schuster, p. 53. Text copyright © 1990 by M. Schuster.

Figure 21.6B: Jones, D. G., Clifford, B. C. (1983) Nature of pathogenicity. In *Cereal Diseases: Their Pathology and Control*, 2nd ed. John Wiley & Sons, New York, pp. 16–26. Reprinted by permission of John Wiley & Sons, Inc.

Figure 21.7: K. Hammond-Kosack, The Sainsbury Laboratory, Norwich, UK; previously unpublished.

Figure 21.8A,B: M. Dow, The Sainsbury Laboratory, Norwich UK; previously unpublished

Figure 21.10A: Franki, R.I.B., Milne, R. G., Hatta, T. (1985) *Atlas of Plant Viruses*, Vol. 2, CRC Press, Boca Raton, FL, p. 138. Copyright © 1985 CRC Press.

Figure 21.10B: Franki, R.I.B., Milne, R. G., Hatta, T. (1985) *Atlas of Plant Viruses*, Vol. 1, CRC Press, Boca Raton, FL, p. 20. Copyright © 1985 CRC Press.

Figure 21.12: Upper panels: Reprinted with permission from Wolf, S., Deom, C. M., Beachy, R. N., Lucas, W. J. (1989) Movement protein of tobacco mosaic virus modifies plasmodesmatal size exclusion limit. *Science* 246: 377–379. Copyright © 1989 American Association for the Advancement of Science. Lower panels: Linstead, P. J., Hills, G. J., Plaskitt, K. A., Willson, I. G., Harker, C. I., Maule, A. J. (1988) The subcellular location

of the gene 1 product of cauliflower mosaic virus is consistent with a function associated with virus spread. *J. Gen. Virol.* 69: 1809–1818. Reprinted with permission of the Society of General Microbiology.

Figure 21.14: Monsanto Co., St. Louis, MO; previously unpublished.

Figure 21.15: Adapted with permission from Lucas, W. J. (1994) Plasmodesmata in relation to viral movement within leaf tissues. *Annu. Rev. Phytopathol.* 32: 387–411. Copyright © 1994 Annual Reviews. www.AnnualReviews.org

Figure 21.16: Gorg, R., Hollricher, K., Schulze-Lefert, P. (1993) Functional analysis and RFLP-mediated mapping of the *Mlg* resistance locus in barley. *Plant J.* 3: 857–866. Used with permission from Blackwell Science Ltd.

Figure 21.17A: Osbourn, A. E. (1999) Antimicrobial phytoprotectants and fungal pathogens: a commentary. *Fungal Genet. Biol.* 26: 163–168. Reprinted by permission of Academic Press. Copyright © 1999.

Figure 21.18B: J. Brown, John Innes Centre, Norwich, UK; previously unpublished.

Figure 21.22: Adapted by permission from Herbers, K., Conrads-Strauch, J., Bonas, U. (1992) Race-specificity of plant resistance to bacterial spot disease determined by repetitive motifs in a bacterial avirulence protein. *Nature* 356: 172–174. Copyright © 1992 Macmillan Magazines Ltd.

Figure 21.23: Three-dimensional model adapted from Vervoort, J., Van Den Hooven, H. W., Berg, A., Vossen, P., Vogel-sand, R., Joosten, M.H.A.J., De Wit, P.G.M. (1997) The race-specific elicitor AVR9 of the tomato pathogen *Cladosporium fulvum*: a cystine knot protein. *FEBS Lett.* 404: 153–158. Copyright © 1997 with permission from Elsevier Science.

Figure 21.27A,B: Reprinted from Kobe, B., Deisenhofer, J. (1995) Proteins with leucine-rich repeats. *Curr. Opin. Struct. Biol.* 5: 409–416. Copyright © 1995 with permission from Elsevier Science.

Figure 21.28B: Carland, F. M., Staskawicz, B. J. (1993) Genetic characterization of the *Pto* locus of tomato: semi-dominance and cosegregation of resistance to *Pseudomonas syringae* pathovar tomato and sensitivity to the insecticide Fenthion. *Mol. Gen. Genet.* 239: 17–27. Copyright © 1993 Springer-Verlag.

Figure 21.28C: Salmeron, J. M., Oldroyd, G.E.D., Rommons, C.M.T., Scofield, S. R., Kim, H.-S., Lavelle, D. T., Dahlbeck, D., Staskawicz, B. J. (1996) Tomato *Prf* is a member of the leucine-rich repeat class of plant disease resistance genes and lies embedded with the *Pto* kinase gene cluster. *Cell* 86: 123–133. Copyright © 1996 Cell Press.

Figure 21.35: Ho, J.-Y., Weide, R., Ma, H. M., van Wordragen, M. F., Lambert, K. N., Koomneef, M., Zabel, P., Williamson, V. M.

(1992) The root-knot nematode resistance gene (*Mf*) in tomato: construction of a molecular linkage map and identification of dominant cDNA markers in resistant genotypes. *Plant J.* 2: 971–973. Used with permission from Elsevier Science.

Figure 21.37B: Bestwick, C. S., Brown, I. R., Bennett, M.H.R., Mansfield, J. W. (1997) Localization of hydrogen peroxide accumulation during the hypersensitive reaction of lettuce cells to *Pseudomonas syringae* pv. *phaseolicola*. *Plant Cell* 9: 209–221.

Figure 21.38: Reprinted by permission from Delledonne, M., Xia, Y., Dixon, R. A., Lamb, C. (1998) Nitric oxide functions as a signal in plant disease resistance. *Nature* 394: 585–588. Copyright © 1998 Macmillan Magazines Ltd.

Figure 21.39: Photograph: Hammond-Kosack, K. E., Jones, J.D.G. (1996) Resistance gene-dependent plant defense responses. *Plant Cell* 8: 1773–1791.

Figure 21.41A: Adapted from Enyedí, A.J., Yalpani, N., Silverman, P., Raskin, I. (1992) Localization, conjugation, and function of salicylic acid in tobacco during the hypersensitive reactants to tobacco mosaic virus. *Proc. Natl. Acad. Sci. USA* 89: 2480–2484.

Figure 21.41B: Hammond-Kosack, K. E., Jones, J.D.G. (1996) Resistance gene-dependent plant defense responses. *Plant Cell* 8: 1773–1791.

Figure 21.42A: Bent, A. F., Innes, R. W., Ecker, J. R., Staskawicz, B. J. (1997) Disease development in ethylene-insensitive *Arabidopsis thaliana* infected with virulent and avirulent *Pseudomonas* and *Xanthomonas* pathogens. *Mol. Plant Microbe Interact.* 5: 372–378. Published by the American Phytopathological Society.

Figure 21.42B: Knoester, M., van Loon, L. C., van den Heuvel, J., Hennig, J., Bol, J. F., Linthorst, H.U.B. (1998) Ethylene-insensitive tobacco lacks nonhost resistance against soil-borne fungi. *Proc. Natl. Acad. Sci. USA* 95: 1933–1937. Copyright © 1998 National Academy of Sciences, U.S.A.

Figure 21.43: Lawton, K. A., Potter, S. L., Uknes, S., Ryals, J. (1994) Acquired resistance signal transduction in *Arabidopsis* is ethylene independent. *Plant Cell* 6: 581–588.

Figure 21.44A,B: Terras, F.R.G., Eggermont, K., Kovaleva, V., Raikhel, N. V., Osborn, R. W., Kester, A., Torrekens, S., Van Leuven, F., Vanderleyden, J., Cammue, V.P.A., Broek-aert, W. F. (1995) Small cyseine-rich antifungal proteins from radish: their role in host defense. *Plant Cell* 7: 573–588.

Figure 21.46: Hammond-Kosack, K. E., Jones, J.D.G. (1996) Resistance gene-dependent plant defense responses. *Plant Cell* 8: 1773–1791.

Figure 21.48: Reprinted from Durner, J., Shah, J., Klessig, D. F. (1997) Salicylic acid and disease resistance in plants. *Trends*

Plant Sci. 2: 266–274. Copyright © 1997 with permission from Elsevier Science.

Figure 21.52: Howe, G. A., Lightner, J., Browse, J., Ryan, C. A. (1996) An octadecanoid pathway mutant (JL5) of tomato is compromised in signaling for defense against insect attack. *Plant Cell* 8: 2067–2077.

Figure 21.54A,B: Lauge, R., Joosten, M.H.A.J., Van den Ackerveken, G.F.J.M., Van den Broek, H.W.J., De Wit, P.J.G.M. (1997) The in planta-produced extracellular proteins ECP1 and ECP2 of *Cladosporium fulvum* are virulence factors. *Mol. Plant Microbe Interact.* 10: 725–734. Reprinted with permission from the American Phytopathological Society.

Figure 21.55A: Adapted with permission from De Wit, P.J.G.M. (1992) molecular characterization of gene-for-gene systems in plant–fungus interactions and the application of avirulence genes in control of plant pathogens. *Annu. Rev. Phytopathol.* 30: 391–418. Copyright © 1992 Annual Reviews. www.AnnualReviews.org

Figure 21.56A: Reprinted by permission from Hain, R., Reif, H.-J., Krause, E., Langebartels, R., Kindl, H., Vornam, B., Wiese, W., Schmelzer, E., Schreier, P. H., Stöcker, R. H., Stenzel, K. (1993) Disease resistance results from foreign phytoalexin expression in a novel plant. *Nature* 361: 153–156. Copyright © 1993 Macmillan Magazines Ltd.

Figure 21.56B: Zhu, Q., Maher, E. A., Masoud, S., Dixon, R. A., Lamb, C. J. (1994) Enhanced protection against fungal attack by constitutive co-expression of chitinase and glucanase genes in transgenic tobacco. *Bio/Technology* 12: 807–812. Used with permission of Nature Publishing Group, New York.

Figure 21.56C: Reprinted with permission from Shah, D. M., Rommens, C.M.T., Beachy, R. N. (1995) Resistance to diseases and insects in transgenic plants: progress and applications to agriculture. *Trends Biotechnol.* 13: 363–368. Copyright © 1995 with permission from Elsevier Science.

Figure 21.56D: Beffa, R. S., Hofer, R.-M., Thomas, M., Meins, F. Jr. (1996) Decreased susceptibility to viral disease of β -1,3-glucanase-deficient plants generated by anti-sense transformation. *Plant Cell* 8: 1001–1011.

Figure 21.57A: de la Fuente-Martinez, J. M., Mosqueda-Cana, G., Alvarez-Morales, A., Herrera-Estrella, L. (1992) Expression of a bacterial phaseolotoxin-resistant ornithyl thanscarbamylyase in transgenic tobacco confers resistance to *Pseudomonas syringae* pv. *phaseolicola*. *Bio/Technology* 10: 905–909. Used with permission of Nature Publishing Group, New York.

Figure 21.57B: Reprinted with permission from Abel, P. P., Nelson, R. S., De, B., Hoffmann, N., Rogers, S. G., Fraley, R. T.,

Beachy, R. N. (1986) Delay of disease development in transgenic plants that express the tobacco mosaic virus coat protein gene. *Science* 232: 738–743. Copyright © 1986 American Association for the Advancement of Science.

Figure 21.57C: Lapidot, M., Gafny, R., Ding, B., Wolf, S., Lucas, W. J., Beachy, R. N. (1993) A dysfunctional movement protein of tobacco mosaic virus that partially modifies the plasmodesmata and limits virus spread in transgenic plants. *Plant J.* 4: 959–970. Used with permission from Blackwell Science Ltd.

Figure 21.58A: Grochulski, P., Masson, L., Borisova, S., Pusztai-Carey, M., Schwartz, J. L., Brousseau, R., Cygler, M. (1995) *Bacillus thuringiensis* CryIA(a) insecticidal toxin: crystal structure and channel formation. *J. Mol. Biol.* 254: 447–464. Reprinted by permission of Academic Press. Copyright © 1999.

Figure 21.58B: Reprinted by permission from Li, J., Carroll, J., Ellar, D. J. (1991) Crystal structure of insecticidal δ -endotoxin from *Bacillus thuringiensis* at 2.5 Å resolution. *Nature* 353: 815–821. Copyright © 1991 Macmillan Magazines Ltd.

Figure 21.59: Photograph: Reprinted with permission from Shah, D. M., Rommens, C.M.T., Beachy, R. N. (1995) Resistance to diseases and insects in transgenic plants: progress and applications to agriculture. *Trends Biotechnol.* 13: 362–368. Copyright © 1995 with permission from Elsevier Science. Graph: Adapted from Munkvold, G. P., Helmut, R. L., Showers, W. B. (1997) Reduced fusarium ear rot and symptomless infection in kernels of maize genetically engineered for European corn borer resistance. *Phytopathology* 87: 1071–1077. Published by the American Phytopathological Society.

Figure 21.60A,B: Reprinted with permission from Tavladoraki, P., Benvenuto, E., Trinca, S., De Martinis, D., Cattaneo, A., Galeffi, P. (1993) Transgenic plants expressing a functional single-chain Fv antibody are specifically protected from virus attack. *Nature* 366: 469–472. Copyright © 1993 Macmillan Magazines Ltd.

Box 21.2: J. Brown, John Innes Centre, Norwich, UK; previously unpublished.

Box 21.3: Adapted from Alfano, J. R., Collmer, A. (1997) The Type III (Hrp) secretion pathway of plant pathogenic bacteria: trafficking harpins, Avr proteins, and death. *J. Bacteriol.* 179: 5655–5662. Used with permission from the American Society for Microbiology.

Box 21.4: Pair of photographs at right: Dinesh-Kumar, S. P., Whitham, S., Choi, D., Hehl, R., Corr, C., Baker, B. (1995) Transposon tagging of tobacco mosaic virus resistance gene *N*: its possible role in the TMV-*N*-mediated signal transduction pathway. *Proc. Natl. Acad. Sci. USA* 92:

4175–4180. Copyright © 1995 National Academy of Sciences, U.S.A.

Box 21.4A–C: Reprinted with permission from Jones, D. A., Thomas, C. M., Hammond-Kosack, K. E., Balint-Kurti, P. J., Jones, J.D.G. (1994) Isolation of the tomato *Cf-9* gene for resistance to *Cladosporium fulvum* by transposon tagging. *Science* 266: 789–793. Copyright © 1994 American Association for the Advancement of Science.

Box 21.5: Adapted from Devoto, A., Piffanelli, P., Nilsson, I., Wallin, E., Panstruga, R., Von Heijne, G., Schulze-Lefert, P. (1999) Topology, subcellular localization, and sequence diversity of the Mlo family in plants. *J. Biol. Chem.* 274: 34993–35004. Copyright © 1999 American Society for Biochemistry and Molecular Biology.

Box 21.7: Whitham, S., McCormick, S., Baker, B. (1996) The *N* gene of tobacco confers resistance to tobacco mosaic virus in transgenic tomato. *Proc. Natl. Acad. Sci. USA* 93: 8776–8781. Copyright © 1996 National Academy of Sciences, U.S.A.

Box 21.8: Jabs, T., Tschoepe, M., Colling, C., Hahlbrock, K., Scheel, D. (1997) Elicitor-stimulated ion fluxes and O₂⁻ from the oxidative burst are essential components in triggering defense gene activation and phytoalexin synthesis in parsley. *Proc. Natl. Acad. Sci. USA* 94: 4800–4805. Copyright © 1997 National Academy of Sciences, U.S.A.

Box 21.9: Photographs: Vernooij, B., Friedrich, L., Morse, A., Reist, R., Kolditz-Jawhar, R., Ward, E., Uknes, S., Kessman, H., Ryals, J. (1994) Salicylic acid is not the translocated signal responsible for inducing systemic acquired resistance but is required in signal transduction. *Plant Cell* 6: 959–965.

Chapter 22

Figure 22.2: L. E. Epple, in Epple, A. O. (1995) *A Field Guide to the Plants of Arizona*. Falcon Press Publishing, Helena, MT.

Figure 22.12: Yamada, S., Katsuhara, M., Kelly, W. B., Michalowski, C. B., Bohnert, H. J. (1995) A family of transcripts encoding water channel proteins: tissue-specific expression in the common ice plant. *Plant Cell* 7: 1129–1142.

Figure 22.13: Adapted from Kononowicz, A. K., Ragothama, K. G., Casa, A. M., Reuveni, M., Watad, A.-E. A., Liu, D., Bressan, R. A., Hasegawa, P. M. (1993) Osmotin: regulation of gene expression and function. In *Plant Response to Cellular Dehydration During Environmental Stress*, T. J. Close and E. A. Bray, eds. American Society of Plant Physiologists, Rockville, MD, pp. 144–158.

Figure 22.17A: V. Ramey, University of Florida Center for Aquatic and Invasive Plants, <http://aquat1.ifas.ufl.edu/hyvepic.htm>. Copyright © 1999 University of Florida.

Figure 22.17B: B. Nelson, <http://aquat1.ifas.ufl.edu/hyvepic.htm>. Copyright ©

1997 Southwest Florida Water Management District.

Figure 22.18: He, C.-J., Morgan, P. W., Drew, M. C. (1996) Transduction of an ethylene signal is required for cell death and lysis in the root cortex of maize during aerenchyma formation induced by hypoxia. *Plant Physiol.* 112: 463–472.

Figure 22.19: Kozłowski, T. T. (1984) *Physiological Ecology*. Academic Press, New York, p. 147. Reprinted by permission of Academic Press. Copyright © 1984.

Figure 22.20: Bowes, B. G. (1996) *A Colour Atlas of Plant Structures*. Manson Publishers, p. 136. Copyright © 1996 Manson Publishers Ltd.

Figure 22.21A: Adapted from Catling, D. (1992) *Rice in Deep Water* (Macmillan, London) and published with permission of the International Rice Research Institute in Kende, H., van der Knaap, E., Cho, H.-T. (1998) Deepwater rice: a model plant to study stem elongation. *Plant Physiol.* 118: 1105–1110.

Figure 22.21B: Kende, H., van der Knaap, E., Cho, H.-T. (1998) Deepwater rice: a model plant to study stem elongation. *Plant Physiol.* 118:1105–1110.

Figure 22.22: Couée, I., Defontaine, S., Carde, J.-P., Pradet, A. (1992) Effects of anoxia on mitochondrial biogenesis in rice shoots: modification of *in organello* translation characteristics. *Plant Physiol.* 98: 411–421.

Figure 22.23: M. S. Martin, University of Illinois at Urbana-Champaign; previously unpublished.

Figure 22.27: Adapted from Roberts, J.K.M., Callis, J., Wemmer, D., Walbot, V., Jardetzky, O. (1984) Mechanism of cytoplasmic pH regulation in hypoxic maize root tips and its role in survival under hypoxia. *Proc. Natl. Acad. Sci. USA* 81: 3379–3383. Used with permission.

Figure 22.29A,B: Adapted from Freeling, M. (1973) Simultaneous induction by anaerobiosis or 2,4-D of multiple enzymes specified by two unlinked genes: differential *Adh1-Adh2* expression in maize. *Mol. Gen. Genet.* 127: 215–227. Copyright © 1973 Springer-Verlag.

Figure 22.30: Adapted from Sachs, M. M., Freeling, M., Okimoto, R. (1980) The anaerobic proteins of maize. *Cell* 20: 761–767. Copyright © 1980 Cell Press.

Figure 22.34A: Reprinted from Jackson, M. (1997) Hormones from roots as signals for the shoots of stressed plants. *Trends Plant Sci.* 2: 22–28. Copyright © 1997 with permission from Elsevier Science.

Figure 22.39: Jacobson, J. S., Hill, A. C., eds. (1970) *Recognition of Air Pollution Injury to Vegetation: A Pictorial Atlas*. Air Pollution Control Association, Pittsburgh, PA.

Figure 22.41A–D: Mansfield, M. A., Key, J. L. (1987) Synthesis of the low molecular weight heat shock proteins in plants. *Plant Physiol.* 84: 1007–1017.

Figure 22.42: Key, J. L., Lin, C. Y., Ceglaz, E., Schoffl, F. (1983) The heat shock response in soybean seedlings. (In *Structure and Function of Plant Genomes*, O. Ciferri and L. Dure, eds.) NATO ASI Ser. A 63: 25–36.

Figure 22.44: Adapted from Morimoto, R. I., Jurivich, D. A., Kroeger, P. E., Mathur, S. K., Murphy, S. P., Nakai, A., Sarge, K., Abravaya, K., Sistonen, L. T. (1994) Regulation of heat shock gene transcription by a family of heat shock factors. In *The Biology of Heat Shock Proteins and Molecular Chaperones*, R. I. Morimoto, A. Tissières, and C. Georgopoulos, eds. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, pp. 417–455. Copyright © 1994 Cold Spring Harbor Laboratory Press.

Box 22.1: Photograph: Xu, D., Duan, X., Wang, B., Hong, B., Ho, T.-H.D., Wu, R. (1996) Expression of a late embryogenesis abundant protein gene, *HVA1*, from barley confers tolerance to water deficit and salt stress in transgenic rice. *Plant Physiol.* 110: 249–257.

Table 22.1: Reprinted with permission from Boyer, J. S. (1982) Plant productivity and environment. *Science* 218: 443–448. Copyright © 1982 American Association for the Advancement of Science.

Chapter 23

Figure 23.1: Ulrich, M.A.E., Allen, W. W. (1992) Strawberry deficiency symptoms: a visual and plant analysis guide to fertilization. Agri. Exp. Station Bull. 1917. Copyright © University of California Board of Regents, Division of Agriculture and Natural Resources, Davis.

Figure 23.3: Adapted from Epstein, E., Rains, D. W., Elzam, O. E. (1962) Resolution of dual mechanisms of potassium absorption by barley roots. *Proc. Natl. Acad. Sci. USA* 49: 684–692. Used with permission.

Figure 23.4: Data from Kochian, L. V., Lucas, W. J. (1982) Potassium transport in corn roots. *Plant Physiol.* 70: 1723–1731.

Figure 23.5: Adapted from Rodriguez-Navarro, A., Blatt, M. R., Slayman, C. L. (1986) A potassium-proton symport in *Neurospora crassa*. *J. Gen. Physiol.* 87: 649–674. By copyright permission from the Rockefeller University Press.

Figure 23.6: Anderson, J. A., Huprikar, S. S., Kochian, L. V., Lucas, W. J., Gaber, R. F. (1992) Functional expression of a probable *Arabidopsis thaliana* potassium channel in *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA* 89: 3736–3740. Used with permission.

Figure 23.7: L. Kochian, Cornell University, Ithaca, NY, and W. Lucas, University of California, Davis; previously unpublished.

Figure 23.9: Wegner, L. H., Raschke, K. (1994) Ion channels in the xylem parenchyma of barley roots. A procedure to isolate protoplasts from this tissue and a patch-clamp exploration of salt passageways into xylem vessels. *Plant Physiol.* 105: 799–813.

Figure 23.10: Reprinted with permission from Schachtman, D. P., Schroeder, J. I. (1994) Structure and transport mechanism of a high-affinity potassium uptake transporter from higher plants. *Nature* 370: 655–658. Copyright © 1994 Macmillan Magazines Ltd.

Figure 23.11: Adapted with permission from Rubio, F., Gassmann, W., Schroeder, J. I. (1995) Sodium-driven potassium uptake by the plant potassium transporter HKT1 and mutations conferring salt tolerance. *Science* 270: 1660–1663. Copyright © 1995 American Association for the Advancement of Science.

Figure 23.12: Adapted from Kim, E. J., Kwak, J. M., Uozumi, N., Schroeder, J. I. (1998) *AtKUP1*: an *Arabidopsis* gene encoding high-affinity potassium transport activity. *Plant Cell* 10: 51–62.

Figure 23.13: Adapted from Kim, E. J., Kwak, J. M., Uozumi, N., Schroeder, J. I. (1998) *AtKUP1*: an *Arabidopsis* gene encoding high-affinity potassium transport activity. *Plant Cell* 10: 51–62.

Figure 23.14: Adapted from Fu, H.-H., Luan, S. (1998) *AtKUP1*: a dual-affinity K⁺ transporter from *Arabidopsis*. *Plant Cell* 10: 63–73.

Figure 23.15: Adapted from Batten, G. D., Wardlaw, I. F. (1987) Senescence and grain development in wheat plants grown with contrasting phosphorus regimes. *Aust. J. Plant Physiol.* 14: 253–268. Reprinted with copyright permission from Commonwealth Scientific and Industrial Research Organization, Melbourne.

Figure 23.16: Dunlop, J., Gardiner, S. (1993) Phosphate uptake, proton extrusion and membrane electropotentials of phosphorus-deficient *Trifolium repens* L. *J. Exp. Bot.* 44: 1801–1808. Used with permission from the Oxford University Press.

Figure 23.17A: D. Redecke, University of Marburg, Germany; previously unpublished.

Figure 23.17B: K. Wex, University of Marburg, Germany; previously unpublished.

Figure 23.18: Johnson, J. F., Vance, C. P., Allen, D. L. (1996) Phosphorus deficiency in *Lupinus albus*. Altered lateral root development and enhanced expression of phosphoenolpyruvate carboxylase. *Plant Physiol.* 112: 31–41.

Figure 23.19A,B: Johnson, J. F., Vance, C. P., Allen, D. L. (1996) Phosphorus deficiency in *Lupinus albus*. Altered lateral root development and enhanced expression of

phosphoenolpyruvate carboxylase. *Plant Physiol.* 112: 31–41.

Figure 23.20: Adapted from Muchhal, U. S., Pardo, J. M., Raghothama, K. G. (1996) Phosphate transporters from the higher plant *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* 93: 10519–10523. Copyright © 1996 National Academy of Sciences, U.S.A.

Figure 23.22: Liu, C., Muchhal, U. S., Uthappa, M., Kononowicz, A. K., Raghothama, K. G. (1998) Tomato phosphate transporter genes are differentially regulated in plant tissues by phosphorus. *Plant Physiol.* 116: 91–99.

Figure 23.24: Welch, R. M., Norvell, W. A., Schaefer, S. C. (1993) Induction of iron(III) and copper(II) reduction in pea roots by Fe and Cu status: does the root-cell plasma-membrane Fe(II)-chelate reductase perform a general role in regulating cation uptake? *Planta* 190: 555–561. Copyright © 1993 Springer-Verlag.

Figure 23.25: Reprinted with permission from Robinson, N. J., Procter, C. M., Connolly, E. L., Guerinot, M. L. (1999) A ferric-chelate reductase for iron uptake from soils. *Nature* 397: 694–697. Copyright © 1999 Macmillan Magazines Ltd.

Figure 23.26: Adapted from Eide, D., Broderius, M., Fett, J., Gurinot, M. L. (1996) A novel iron-regulated metal transporter from plants identified by functional expression in yeast. *Proc. Natl. Acad. Sci. USA* 93: 5624–5628. Copyright © 1996 National Academy of Sciences, U.S.A.

Figure 23.28: C. K. Cohen and L. V. Kochian, Cornell University, Ithaca, NY; previously unpublished.

Figure 23.29: Ma, J. F., Nomoto, K. (1993) Two related biosynthetic pathways of mugenic acids in gramineous plants. *Plant Physiol.* 102: 373–378.

Figure 23.31: Adapted from Delhaize, E., Ryan, P. R., Randall, P. J. (1993) Aluminum tolerance in wheat (*Triticum aestivum* L.). *Plant Physiol.* 103: 695–702.

Figure 23.32: Lasat, M. M., Baker, A.J.M., Kochian, L. V. (1996) Physiological characterization of root Zn²⁺ absorption and translocation to shoots in Zn hyperaccumulator and nonaccumulator species of *Thlaspi*. *Plant Physiol.* 112: 1715–1722.

Figure 23.33: I. Raskin, Rutgers University, NJ; previously unpublished.

Figure 23.34: Martin, R. B. (1988) Bioinorganic chemistry of aluminum. In *Metal Ions in Biological Systems: Aluminum and Its Role in Biology*, Vol. 24, H. Sigel and A. Sigel, eds. Marcel Dekker, New York, pp. 2–57. Reprinted by courtesy of Marcel Dekker, Inc.

Figure 23.35: Adapted from Delhaize, E., Ryan, P. R., Randall, P. J. (1993) Aluminum tolerance in wheat (*Triticum aestivum* L.). *Plant Physiol.* 103: 695–702.

Box 23.2: Adapted from Nissen, P. (1989)

Multiphasic uptake of potassium by corn roots. *Plant Physiol.* 89: 231–237.

Box 23.3: Photograph: L. V. Kochian, Cornell University, Ithaca, NY; previously unpublished.

Table 23.1: Epstein, E. (1972) *Mineral Nutrition of Plants: Principles and Perspectives*. John Wiley & Sons, New York. Reprinted by permission of John Wiley & Sons, Inc.

Chapter 24

Figure 24.3A: R. Croteau, Washington State University, Pullman; previously unpublished.

Figure 24.3B–D: Wise, M. L., Croteau, R. (1999) Biosynthesis of monoterpenes. In *Comprehensive Natural Products Chemistry, Vol. 2. Isoprenoids Including Carotenoids and Steroids*, D. E. Cane, ed. Pergamon/Elsevier, Oxford, UK, pp. 97–153. Copyright © 1999 with permission from Elsevier Science.

Figure 24.5: Adapted from McCaskill, D., Croteau, R. (1998) Some caveats for bioengineering terpenoid metabolism in plants. *Trends Biotechnol.* 16: 349–354. Copyright © 1998 with permission from Elsevier Science.

Figure 24.11: Reprinted with permission from Gijzen, M., Lewinsohn, E., Savage, T. J., Croteau, R. B. (1993) Conifer monoterpenes: biochemistry and bark beetle chemical ecology. *ACS Symp. Ser.* 525: 8–22. Copyright © 1993 American Chemical Society.

Figure 24.12: Reprinted with permission from Starks, C. M., Back, K., Chappell, J., Noel, J. P. (1997) Structural basis for cyclic terpene biosynthesis by 5-*epi*-aristolochene synthase from tobacco. *Science* 277: 1815–1820. Copyright © 1997 American Association for the Advancement of Science.

Figure 24.18A: T. M. Kutchan, Leibniz Institut für Pflanzenbiochemie, Halle, Germany; previously unpublished.

Figure 24.18B: Photograph courtesy of the Ministry of Culture Archaeological Receipts Fund, Athens, Greece.

Figure 24.19B: The Metropolitan Museum of Art, New York, NY.

Figure 24.21B: V. Rickl, Universität München, Germany; previously unpublished.

Figure 24.30: M. H. Zenk, Universität München, Germany; previously unpublished.

Figure 24.36: M. H. Zenk, Universität München, Germany; previously unpublished.

Figure 24.47D: L. B. Davin, Washington State University, Pullman; previously unpublished.

Figure 24.48A: L. B. Davin, Washington State University, Pullman; previously unpublished.

Figure 24.48B: Hemp-Museum, Berlin, Germany; previously unpublished.

Figure 24.55: Photograph: L. B. Davin, Washington State University, Pullman; previously unpublished.

Figure 24.57A,C: L. B. Davin, Washington State University, Pullman; previously unpublished.

Figure 24.57B: Photograph: G.H.N. Towers, R. A. Norton, University of British Columbia, Vancouver, Canada; previously unpublished.

Figure 24.59: Data from Adler, E. (1977) Lignin chemistry: past, present and future. *Wood Sci. Technol.* 11: 169–218, and Brunow, G., Kilpeläinen, I., Sipilä, J., Syrjänen, K., Karhunen, P., Setälä, H., Rummakko, P. (1998) Oxidative coupling of phenols and the biosynthesis of lignin. *ACS Symp. Ser.* 697: 131–147.

Figure 24.60B: L. B. Davin, L. H. Levine, Washington State University, Pullman; previously unpublished.

Figure 24.61A–C: L. B. Davin, L. H. Levine, Washington State University, Pullman; previously unpublished.

Figure 24.62: L. H. Levine, Washington State University, Pullman; previously unpublished.

Figure 24.65: Photograph: M. A. Bernards, University of Western Ontario, London, Canada; previously unpublished

Figure 24.67: Photograph: L. B. Davin, Washington State University, Pullman; previously unpublished.

Figure 24.68: Photograph: T. Lumpkin, US Department of Agriculture, Washington State University, Pullman; previously unpublished.

Figure 24.69: Photographs: L. B. Davin (alfalfa plant), M. L. Kahn (nodules), Washington State University, Pullman; previously unpublished.

Figure 24.73: Photographs: G.H.N. Towers, University of British Columbia, Vancouver, Canada; previously unpublished.

Figure 24.77: Photograph: L. B. Davin, Washington State University, Pullman; previously unpublished.

Box 24.2A: French Pharmacopée Royale; photograph: T. Kutchan, Leibniz Institut für Pflanzenbiochemie, Halle, Germany; previously unpublished.

Box 24.2B: Residenz Pharmacy, Munich, Germany; photograph: T. Kutchan, Leibniz Institut für Pflanzenbiochemie, Halle, Germany; previously unpublished.

Box 24.3A,B: T. Kutchan, Leibniz Institut für Pflanzenbiochemie, Halle, Germany; previously unpublished.

Box 24.4: Diagram adapted from Setchell, K.C.R., Lawson, A. M., Borriello, S. P., Adlercreutz, H., Axelson, M. (1982) Formation of lignan by intestinal microflora. *Falk Symp.* 31: 93–97. Used with kind permission from Kluwer Academic Publishers.

Box 24.5A: Gang, D. R., Fujita, M., Davin, L. B., Lewis, N. G. (1998) The “abnormal lignins”: mapping heartwood formation

through the lignan biosynthetic pathway. In *Lignan and Lignan Biosynthesis (ACS Symp. Ser.)*, N. G. Lewis, and S. Sarkanen, eds., Washington, DC. Copyright © 1998 American Chemical Society.

Box 24.5B,C: L. Shain, W. E. Hillis, Commonwealth Scientific and Industrial Re-

search Organization, Clayton South MDC, Victoria, Australia; previously unpublished.

Box 24.5D: Adapted from Chattaway, M. M. (1949) The development of tyloses and secretion of gum in heartwood formation. *Aust. J. Sci. Res.* 227-240.

Box 24.6: L. B. Davin, Washington State University, Pullman; previously unpublished.